

An Estimation of the Length of a Convex Curve in Two-Dimensional Aleksandrov Spaces

Alexander A. Borisenko

In the paper, a generalization of the Toponogov theorem about the length of a curve in a two-dimensional Riemannian manifold is proved for the case of two-dimensional Aleksandrov spaces.

Key words: λ -convex curve, two-dimensional Aleksandrov space.

Mathematical Subject Classification 2010: 53C44, 52A40

Let R be an Aleksandrov space of curvature $\geq c$ homeomorphic to a disc (see [1, p. 308]). Suppose G is a domain in R that is bounded by a rectifiable curve γ . Denote by $\tau(\gamma_1)$ the integral geodesic curvature (the swerve) of a subarc γ_1 of γ [1, p. 309]. The curve γ is called λ -convex with $\lambda > 0$ if any subarc γ_1 of γ satisfies

$$\frac{\tau(\gamma_1)}{s(\gamma_1)} \geq \lambda > 0, \quad (1)$$

where $s(\gamma_1)$ is the length of γ_1 . For regular curves in a two-dimensional Riemannian manifold this condition is equivalent to the assumption that the geodesic curvature at each point of this curve is $\geq \lambda > 0$. In the general case the condition (1) allows γ to have corner points.

We prove the following theorem.

Theorem 1. *Let G be a domain homeomorphic to a disc and G lie in a two-dimensional Aleksandrov space of curvature $\geq c$ (in the sense of Aleksandrov).*

I. *If the boundary curve γ of G is λ -convex and $c + \lambda^2 > 0$, then the length $s(\gamma)$ of γ satisfies*

1. $s(\gamma) \leq \frac{2\pi}{\lambda}$ for $c = 0$;
2. $s(\gamma) \leq \frac{2\pi\sqrt{c}}{\sqrt{c + \lambda^2}}$ for $c > 0$;
3. $s(\gamma) \leq \frac{2\pi\sqrt{-c}}{\sqrt{c + \lambda^2}}$ for $c < 0$.

- II. *All these inequalities become equalities if and only if the domain G is a disc on the plane of constant curvature c .*

This theorem is a generalization of the Toponogov theorem [10] about the length of a curve in a two-dimensional Riemannian manifold. We need the following statements to prove Theorem 1.

Theorem A (A.D. Aleksandrov [1, p. 269]). *A metric space with intrinsic metric of curvature $\geq c$ homeomorphic to a sphere is isometric to a closed convex surface in a simply connected space of constant curvature c .*

Theorem B (A.V. Pogorelov [9, pp. 119–167, 267, 320–321], [4]). *Closed isometric convex surfaces in the three-dimensional Euclidean and spherical spaces are equal up to a rigid motion.*

Theorem C (A.D. Milka [8]). *Closed isometric convex surfaces in the three-dimensional Lobachevsky space are equal up to a rigid motion.*

Theorem D (W. Blaschke [2]). *Let γ be a closed embedded C^2 regular curve in the Euclidean plane.*

- I. *If the curvature k of γ at each its point P satisfies*

$$k \geq \lambda > 0,$$

then the curve belongs to the disc that is bounded by the circle of radius $R = 1/\lambda$ tangent to the curve at P .

- II. *If the curvature k of γ at each its point P satisfies*

$$0 \leq k \leq \lambda,$$

then the circle of radius $R = 1/\lambda$ tangent to the curve at P belongs to the domain G that is bounded by the curve γ .

Theorem D remains true if the condition for the curvature k of γ is replaced by the same condition for the specific curvature $\frac{\tau(\gamma_1)}{s(\gamma_1)}$ for any arc.

Lemma 1. *Let γ be a closed embedded rectifiable curve in the Euclidean plane.*

- I. *If for any subarc γ_1 of γ the specific curvature $\frac{\tau(\gamma_1)}{s(\gamma_1)}$ satisfies*

$$\frac{\tau(\gamma_1)}{s(\gamma_1)} \geq \lambda > 0,$$

then the curve γ belongs to the disc that is bounded by the circle of radius $R = 1/\lambda$ tangent to the support straight line of γ at its point P .

II. If for any subarc γ_1 of γ the specific curvature $\frac{\tau(\gamma_1)}{s(\gamma_1)}$ satisfies

$$0 \leq \frac{\tau(\gamma_1)}{s(\gamma_1)} \leq \lambda,$$

then the circle of radius $R = 1/\lambda$ tangent to the curve at its point P belongs to the domain G that is bounded by the curve γ .

Proof. I. In this case the support function $h(\phi)$, $0 \leq \phi \leq 2\pi$, of the curve γ is $C^{1,1}$ regular and a.e. it satisfies the equation

$$h + h'' = R, \quad 0 \leq R \leq \frac{1}{\lambda},$$

where R is the radius of curvature of γ . Therefore,

$$h(\phi) = \int_0^\phi R(\sigma) \sin(\phi - \sigma) d\sigma,$$

and the proof coincides with Blaschke's proof [2].

II. The position vector $r(s)$ of the curve γ is a $C^{1,1}$ regular vector function. Fix an initial point P_0 on γ and denote by e_1 the unit tangent vector of γ at P_0 , and by e_2 the unit normal vector of γ at P_0 . Let $P(s)$ be a point on γ such that the length of the arc $\gamma(s) = P_0P(s)$ equals s . The function $\tau(s) = \tau(\gamma(s))$ is the integral geodesic curvature of the arc $\gamma(s)$, and $\tau(s) \leq \lambda s$. Therefore,

$$r'(s) = \cos \tau(s) e_1 + \sin \tau(s) e_2. \tag{2}$$

If we compare (2) with the equation for the circle of radius $1/\lambda$, we obtain the proof. □

H. Karcher proved a generalization of the Blaschke theorem for regular curves in the spherical space \mathbb{S}^2 and in the Lobachevsky space \mathbb{H}^2 [7]. We formulate Lemma 2 for the cases where the curvature of \mathbb{S}^2 is equal to 1 and the curvature of \mathbb{H}^2 is equal to -1 . Lemma 2 remains true for the planes of any constant curvature c and the proof is the same.

Lemma 2. *Let γ be a closed embedded rectifiable curve in \mathbb{H}^2 or \mathbb{S}^2 .*

I. *If the specific curvature satisfies*

$$\frac{\tau(\gamma_1)}{s(\gamma_1)} \geq \coth R_0 = \lambda$$

for any subarc γ_1 of γ in \mathbb{H}^2 , then the curve γ belongs to the disc that is bounded by the circle of radius R_0 tangent to the support straight line of γ at a point $P \in \gamma$.

II. If the specific curvature satisfies

$$\frac{\tau(\gamma_1)}{s(\gamma_1)} \geq \cot R_0 = \lambda,$$

for any subarc γ_1 of γ in \mathbb{S}^2 , then the curve γ belongs to the disc that is bounded by the circle of radius R_0 tangent to the support straight line of γ at a point $P \in \gamma$.

Proof. The curve γ is a closed convex curve. At any point P of γ there exists a support straight line (a geodesic line in the plane of constant curvature).

I: $\gamma \in \mathbb{H}^2$. Let S be a circle of radius R_0 tangent to the support straight line of γ from the side containing γ . Assume that the center of the circle S is the origin of the coordinate system in the Cayley–Klein model of the Lobachevsky plane and it is also the origin for the support function h of the curve γ . The support function h is $C^{1,1}$ regular and a.e. the radius of curvature R of γ equals

$$R = \frac{g + g''}{\left(1 - \frac{(g')^2}{1+g^2}\right)^{3/2}}, \quad (3)$$

where $g(h) = \tanh h$ is the support function for the curve $\tilde{\gamma}$, and $\tilde{\gamma}$ is the image of γ under the geodesic map from \mathbb{H}^2 to \mathbb{E}^2 [5, 6]. The radius of curvature \tilde{R} of $\tilde{\gamma}$ is a.e. equal to

$$\tilde{R} = R \left(1 - \frac{(g')^2}{1+g^2}\right)^{3/2}, \quad 0 \leq \tilde{R} \leq R. \quad (4)$$

The image of the circle S under the geodesic map is the circle \tilde{S} in the Euclidean plane \mathbb{E}^2 with the center at the origin of a Cartesian orthogonal coordinate system. The curvature of \tilde{S} equals $\coth R_0$. From Lemma 1(I), it follows that $\tilde{\gamma}$ belongs to the disc bounded by the circle \tilde{S} . Applying the inverse geodesic transformation, we obtain that the curve γ belongs to the disc that is bounded by the circle S in the Lobachevsky plane \mathbb{H}^2 .

II: $\gamma \in \mathbb{S}^2$. Let $\bar{\gamma}$ be the polar to the curve γ in \mathbb{S}^2 . The position vector of $\bar{\gamma}$ is $C^{1,1}$ regular and its curvature is $\leq \tan R_0$ a.e. Let P_0 be a point on γ and \bar{S} be a circle of radius $\pi/2 - R_0$ tangent to $\bar{\gamma}$ at the point \bar{P}_0 . The curvature of this circle is equal to $\tan R_0$. The center \bar{O} of \bar{S} is the south pole of the sphere. Consider the geodesic map of the sphere \mathbb{S}^2 into the plane tangent to \mathbb{S}^2 at the point \bar{O} . The curve $\bar{\gamma}$ is mapped to a curve $\tilde{\bar{\gamma}} \in \mathbb{E}^2$, and the circle \bar{S} is mapped to a circle $\tilde{\bar{S}}$ of curvature $\tan R_0$. The curvatures satisfy $\tilde{\bar{k}}(\tilde{\bar{\gamma}}) \leq \bar{k}(\bar{\gamma}) \leq \tan R_0$. From Lemma 1(II) it follows that the circle $\tilde{\bar{S}}$ belongs to the domain that is bounded by the curve $\tilde{\bar{\gamma}}$. Applying the inverse geodesic transformation, we obtain that the circle \bar{S} belongs to the domain bounded by $\bar{\gamma}$ and the polar curve γ belongs to the disc bounded by the polar circle S of radius R_0 . \square

Proof of Theorem 1. Let G_1 and G_2 be two copies of the domain G . Let us glue the domains G_1 and G_2 along their boundary curves γ_1 and γ_2 by an

isometry between these curves. We obtain a manifold F homeomorphic to the two-dimensional sphere with an intrinsic metric. Since the sum of the integral geodesic curvatures of any two identified arcs of the boundary curves is non-negative, from the Aleksandrov gluing theorem [1, p. 318] it follows that F is an Aleksandrov space of curvature $\geq c$. By Theorem A, this manifold can be isometrically embedded as a closed convex surface F_1 in the simply-connected space $M^3(c)$ of constant curvature c . From Theorem B and C it follows that this surface is unique up to a rigid motion.

By *plane domains* we will understand domains on totally geodesic two-dimensional surfaces in spaces of constant curvature; similarly we will call geodesic lines in these spaces as *lines*.

Perform the reflection of the surface F_1 with respect to a plane π passing through three points on γ that do not belong to any line. We will get a mirrored surface F_2 . The domains G_1 and G_2 are mapped to domains \tilde{G}_1 and \tilde{G}_2 on F_2 ; the curve γ is mapped to $\tilde{\gamma}$. But G_1 is isometric to G_2 and \tilde{G}_2 is isometric to \tilde{G}_1 . Let us reverse the orientation of the domains \tilde{G}_1, \tilde{G}_2 . Then the surface F_2 will be isometric to F_1 and they will have the same orientation. By Theorems B and C, the surface F_1 can be mapped to the surface F_2 by a rigid motion of the ambient space. But the three points of the curve γ are fixed under this rigid motion. Thus it follows that this motion is the identity mapping and, moreover, the curve γ coincides with the curve $\tilde{\gamma}$. Such a situation is possible only when the curve γ is a plane curve and it is the boundary of a convex cup isometric to the domain G . Recall that the *convex cup* is a convex surface with a planar boundary curve γ such that the surface is a graph over a plane domain \bar{G} enclosed by γ . Note that since γ is a convex curve on the plane, the integral geodesic curvature of any arc of the curve γ is non-negative if γ is viewed as a curve on the cup and as a curve on a plane [3].

Let us show that the integral geodesic curvature of any arc of γ calculated on G is not less than its corresponding integral geodesic curvature calculated on the cup G . This means that γ as the boundary curve of \bar{G} is also λ -convex.

Recall that the intrinsic curvature $\omega(D)$ of a Borel set D on a convex surface in a space of constant curvature c is

$$\omega(D) = \psi(D) + cF(D),$$

where $\psi(D)$ is the extrinsic curvature, $F(D)$ is the area of D [1, p. 397]. Consider a closed convex surface M bounded by G and the plane domain \bar{G} , and a surface \bar{M} composed of the double-covered domain \bar{G} .

The intrinsic curvature concentrated on γ equals

$$\omega(\gamma) = \tau_\gamma(G) + \tau_\gamma(\bar{G}),$$

where $\tau_\gamma(G), \tau_\gamma(\bar{G})$ are the integral geodesic curvatures of γ computed in G and \bar{G} respectively.

Since $F(\gamma) = 0$, we have

$$\psi_M(\gamma) = \tau_\gamma(G) + \tau_\gamma(\bar{G}),$$

$$\psi_{\overline{M}}(\gamma) = 2\tau_\gamma(\overline{G}).$$

From the definition of the extrinsic curvature [1, p. 398] it follows that $\psi_{\overline{M}}(\gamma) \geq \psi_M(\gamma)$ because each plane supporting M at a point of γ is also supporting \overline{M} . Thus we obtain $\tau_\gamma(\overline{G}) \geq \tau_\gamma(G)$. Moreover, this inequality holds for any subarc of γ .

I. The curve γ is a λ -convex curve lying in the plane of constant curvature c . From Lemmas 1 and 2, it follows that the curve γ belongs to a disc bounded by the circle of radius R_0 . The curvature and the length s of this circle equal

1. $\lambda = \frac{1}{R_0}$, $s = 2\pi R_0$ for $c = 0$;
2. $\lambda = \sqrt{c} \cot \sqrt{c} R_0$, $s = 2\pi \sin \sqrt{c} R_0$ for $c > 0$;
3. $\lambda = \sqrt{-c} \coth \sqrt{-c} R_0$, $s = 2\pi \sinh \sqrt{-c} R_0$ for $c < 0$.

The curve γ on the plane of constant curvature c bounds the convex domain G . It follows that the length of γ satisfies

$$s(\gamma) \leq \begin{cases} \frac{2\pi}{\lambda} & \text{if } c = 0 \\ \frac{\lambda}{2\pi\sqrt{c}} & \text{if } c > 0 \\ \frac{\sqrt{c + \lambda^2}}{2\pi\sqrt{-c}} & \text{if } c < 0. \end{cases} \quad (5)$$

II. Suppose that there is equality in (5). Then the domain \overline{G} is a disc bounded by the circle γ . Furthermore, $\tau_\gamma(\overline{G}) = \tau_\gamma(G)$ and the intrinsic curvature of γ satisfies $\omega_M(\gamma) = \omega_{\overline{M}}(\gamma) = 2\tau_\gamma(\overline{G})$, and the extrinsic curvature for any subarc γ_1 of γ satisfies

$$\psi_M(\gamma) = \psi_{\overline{M}}(\gamma). \quad (6)$$

It follows that the surfaces M and \overline{M} coincide, M is a double-covered disk and thus G is a disk. If M does not coincide with \overline{M} , then there exists a set of a positive measure of supporting planes to \overline{M} along γ which are not supporting planes to M . It follows that the extrinsic curvatures of M and \overline{M} along γ do not coincide. This contradicts equality (6), and thus Theorem 1 is proved. \square

References

- [1] A.D. Aleksandrov, *A.D. Aleksandrov selected works, II, Intrinsic geometry of convex surfaces*, (Edited by S. S. Kutateladze; Translated from Russian by S. Vakhrameyev), Chapman & Hall/CRC, Boca Raton, FL, 2006.
- [2] W. Blaschke, *Kreis und Kugel*, Walter de Gruyter & Co., Berlin, 1956 (German).
- [3] A.A. Borisenko, *Reverse isoperimetric inequality in two-dimensional Aleksandrov spaces*, Proc. Amer. Math. Soc. **145** (2017), 4465–4471.
- [4] A.A. Borisenko, *About Kharkiv Mathematicians, Members of the Academy of Sciences of Ukraine*, Zh. Mat. Fiz. Anal. Geome. **15** (2019), No. 2, 288–303 (Russian).
- [5] A.A. Borisenko and K.D. Drach, *Extreme properties of curves with bounded curvature on a sphere*, J. Dyn. Control Syst. **21** (2015), No. 3, 311–327.

- [6] K.D. Drach, *About the isoperimetric property of λ -convex lunes on the Lobachevsky plane*, Dopov. Nats. Akad. Nauk Ukr. **11** (2014), 11–15 (Russian); Engl. transl.: preprint, <https://arxiv.org/abs/1402.2688>.
- [7] H. Karcher, *Umkreise und Inkreise konvexer Kurven in der spharischen und der hyperbolischen Geometrie*, Math. Ann. **177** (1968), 122–132 (German).
- [8] A.D. Milka, *Unique determinacy of general closed convex surfaces in Lobachevsky space*, Ukrain. Geom. Sb. **23** (1980), 99–107 (Russian).
- [9] A.V. Pogorelov, *Extrinsic Geometry of Convex Surfaces*, Translations of Mathematical Monographs, **35**, Amer. Math. Soc, Providence, R.I., 1973.
- [10] V. A. Toponogov, *Estimation of the length of a convex curve on a two-dimensional surface*, Sibirsk. Mat. Zh. **4** (1963), No. 5, 1189–1183 (Russian).

Received June 2, 2020.

Alexander A. Borisenko,

B. Verkin Institute for Low Temperature Physics and Engineering of the National Academy of Sciences of Ukraine, 47 Nauky Ave., Kharkiv, 61103, Ukraine,

E-mail: aborisenk@gmail.com

Про оцінку довжини опуклої кривої у двовимірному просторі Александрова

Alexander A. Borisenko

Доведено узагальнення теореми Топоногова про довжину кривої у двовимірному рімановому просторі на випадок двовимірного простору Александрова.

Ключові слова: λ -опуклі криві, двовимірний простір Александрова