

# Dissipative Extensions of Linear Relations Generated by Integral Equations with Operator Measures

Vladislav M. Bruk

In the paper, a minimal relation  $L_0$  generated by an integral equation with operator measures is defined and a description of the adjoint relation  $L_0^*$  is given. For this minimal relation, we construct a space of boundary values (a boundary triplet) satisfying the abstract "Green formula" and get a description of maximal dissipative (accumulative) and also self-adjoint extensions of the minimal relation.

*Key words:* Hilbert space, linear relation, integral equation, dissipative extension, self-adjoint extension, boundary value, operator measure

*Mathematical Subject Classification 2010:* 47A10, 46G12, 45N05

## 1. Introduction

In the study of linear operators and relations generated by differential or integral equations with boundary conditions there often arises a problem of finding the boundary conditions that determine an operator or a relation with preassigned properties. A classical example of the solution to this problem is the description of self-adjoint extensions of a symmetric operator generated by an ordinary differential expression. The description was given by M.G. Krein in [17] (see also [18, Chap. 5]).

The method proposed by M. G. Krein essentially uses the finite dimensionality of defect subspaces of the symmetric operator. Therefore it is difficult to apply the results obtained in [17] to operators with infinite defect indices. A significant advance in overcoming these difficulties was made by F. S. Rofe-Beketov [20], who was the first to use linear relations for describing self-adjoint extensions of the minimal operator generated by a differential expression with bounded operator coefficients. The results obtained in [20] were later generalized both to the case of more general (accumulative and dissipative) extensions [14] and to the case of differential expressions with unbounded operator coefficients (see monographs [13] and [21] for detailed bibliography).

In this paper we consider the integral equation

$$y(t) = x_0 - iJ \int_{[a,t)} d\mathbf{p}(s)y(s) - iJ \int_{[a,t)} d\mathbf{m}(s)f(s), \quad (1.1)$$

where  $y$  is an unknown function,  $a \leq t \leq b$ ;  $J$  is an operator in a separable Hilbert space  $H$ ,  $J = J^*$ ,  $J^2 = E$  ( $E$  is an identical operator);  $\mathbf{p}$ ,  $\mathbf{m}$  are the operator-valued measures defined on Borel sets  $\Delta \subset [a, b]$  that take values in the set of linear bounded operators acting in  $H$ ;  $x_0 \in H$ ,  $f \in L_2(H, d\mathbf{m}; a, b)$ . We assume that the measures  $\mathbf{p}$ ,  $\mathbf{m}$  have bounded variations,  $\mathbf{p}$  is self-adjoint and  $\mathbf{m}$  is non-negative.

We define a minimal relation  $L_0$  generated by equation (1.1) and give a description of the adjoint relation  $L_0^*$ . For this minimal relation, we construct a space of boundary values (boundary triplet) satisfying the abstract ‘‘Green formula’’ (see [4, 5, 16]) and get a description of maximal dissipative (accumulative) and also self-adjoint extensions of the minimal relation.

If the measures  $\mathbf{p}$ ,  $\mathbf{m}$  are absolutely continuous (i.e.,  $\mathbf{p}(\Delta) = \int_{\Delta} p(t) dt$ ,  $\mathbf{m}(\Delta) = \int_{\Delta} m(t) dt$  for all Borel sets  $\Delta \subset [a, b]$ , where the functions  $\|p(t)\|$ ,  $\|m(t)\|$  belong to  $L_1(a, b)$ ), then integral equation (1.1) is transformed into a differential equation with a non-negative weight operator function. Linear relations and operators generated by such differential equations were considered in many works (see [6, 7, 19], further detailed bibliography can be found, for example, in [3, 15]).

The study of integral equation (1.1) differs essentially from the study of differential equations by the presence of the following features:

- i) a representation of a solution of equation (1.1) using an evolutionary family of operators is possible if the measures  $\mathbf{p}$ ,  $\mathbf{m}$  do not have common single-point atoms (see [8]);
- ii) the Lagrange formula contains summands that are related to single-point atoms of the measures  $\mathbf{p}$ ,  $\mathbf{m}$  (see [9]).

Note that this paper partially corrects the errors made in [10].

Under tighter assumptions imposed on the measures  $\mathbf{p}$ ,  $\mathbf{m}$ , a description of self-adjoint or maximal dissipative (accumulative) extension of  $L_0$  is given in the papers: [9] (where  $\mathbf{m}$  is the usual Lebesgue measure on  $[a, b]$  and the measure  $\mathbf{p}$  has a finite number of single-point atoms); [11] (where  $\mathbf{m}$  is the usual Lebesgue measure on  $[a, b]$  and the set of single-point atoms of the measure  $\mathbf{p}$  can be arranged as an increasing sequence converging to  $b$ ); [12] (where  $\mathbf{m}$  is a non-negative continuous measure and the measure  $\mathbf{p}$  is the same as in [11]). In [9, 11],  $L_0$ ,  $L_0^*$  are operators.

## 2. Preliminary assertions

Let  $H$  be a separable Hilbert space with a scalar product  $(\cdot, \cdot)$  and a norm  $\|\cdot\|$ . We consider a function  $\Delta \rightarrow \mathbf{P}(\Delta)$  defined on Borel sets  $\Delta \subset [a, b]$  that takes values in the set of linear bounded operators acting in  $H$ . The function  $\mathbf{P}$  is called an operator measure on  $[a, b]$  (see, for example, [2, Chap. 5]) if it is zero on the empty set and the equality

$$\mathbf{P} \left( \bigcup_{n=1}^{\infty} \Delta_n \right) = \sum_{n=1}^{\infty} \mathbf{P}(\Delta_n)$$

holds for disjoint Borel sets  $\Delta_n$ , where the series converges weakly. Further, we extend any measure  $\mathbf{P}$  on  $[a, b]$  to a segment  $[a, b_0]$  ( $b_0 > b$ ) letting  $\mathbf{P}(\Delta) = 0$  for each Borel set  $\Delta \subset (b, b_0]$ .

By  $\mathbf{V}_\Delta(\mathbf{P})$ , we denote

$$\mathbf{V}_\Delta(\mathbf{P}) = \rho_{\mathbf{P}}(\Delta) = \sup \sum_n \|\mathbf{P}(\Delta_n)\|,$$

where the supremum is taken over finite sums of disjoint Borel sets  $\Delta_n \subset \Delta$ . The number  $\mathbf{V}_\Delta(\mathbf{P})$  is called the variation of the measure  $\mathbf{P}$  on the Borel set  $\Delta$ . Suppose that the measure  $\mathbf{P}$  has the bounded variation on  $[a, b]$ . Then for  $\rho_{\mathbf{P}}$ -almost all  $\xi \in [a, b]$  there exists an operator function  $\xi \rightarrow \Psi_{\mathbf{P}}(\xi)$  such that  $\Psi_{\mathbf{P}}$  possesses the values in the set of linear bounded operators acting in  $H$ ,  $\|\Psi_{\mathbf{P}}(\xi)\| = 1$ , and the equality

$$\mathbf{P}(\Delta) = \int_{\Delta} \Psi_{\mathbf{P}}(s) d\rho_{\mathbf{P}} \tag{2.1}$$

holds for each Borel set  $\Delta \subset [a, b]$ . The function  $\Psi_{\mathbf{P}}$  is uniquely determined up to values on a set of zero  $\rho_{\mathbf{P}}$ -measure. Integral (2.1) converges in the sense of usual operator norm ([2, Chap. 5]).

Further,  $\int_{t_0}^t$  stands for  $\int_{[t_0, t]}$  if  $t_0 < t$ , for  $-\int_{[t, t_0]}$  if  $t_0 > t$ , and for 0 if  $t_0 = t$ . A function  $h$  is integrable with respect to the measure  $\mathbf{P}$  on a set  $\Delta$  if there exists the Bochner integral

$$\int_{\Delta} \Psi_{\mathbf{P}}(t)h(t) d\rho_{\mathbf{P}} = \int_{\Delta} (d\mathbf{P}) h(t).$$

Then the function

$$(y(t) = \int_{t_0}^t (d\mathbf{P}) h(s)$$

is continuous from the left.

By  $\mathcal{S}_{\mathbf{P}}$ , denote a set of single-point atoms of the measure  $\mathbf{P}$  (i.e., a set  $t \in [a, b]$  such that  $\mathbf{P}(\{t\}) \neq 0$ ). The set  $\mathcal{S}_{\mathbf{P}}$  is at most countable. The measure  $\mathbf{P}$  is continuous if  $\mathcal{S}_{\mathbf{P}} = \emptyset$ , it is self-adjoint if  $(\mathbf{P}(\Delta))^* = \mathbf{P}(\Delta)$  for each Borel set  $\Delta \subset [a, b]$ , it is non-negative if  $(\mathbf{P}(\Delta)x, x) \geq 0$  for all Borel sets  $\Delta \subset [a, b]$  and for all elements  $x \in H$ .

In Lemma 2.1 below,  $\mathbf{p}_1, \mathbf{p}_2, \mathbf{q}$  are operator measures having bounded variations and taking values in the set of linear bounded operators acting in  $H$ . Suppose that the measure  $\mathbf{q}$  is self-adjoint and assume that these measures are extended on the segment  $[a, b_0] \supset [a, b_0] \supset [a, b]$  in the manner described above.

**Lemma 2.1** ([9]). *Let  $f, g$  be functions integrable on  $[a, b_0]$  with respect to the measure  $\mathbf{q}$  and  $y_0, z_0 \in H$ . Then the functions*

$$\begin{aligned} y(t) &= y_0 - iJ \int_{t_0}^t d\mathbf{p}_1(s)y(s) - iJ \int_{t_0}^t d\mathbf{q}(s)f(s), \\ z(t) &= z_0 - iJ \int_{t_0}^t d\mathbf{p}_2(s)z(s) - iJ \int_{t_0}^t d\mathbf{q}(s)g(s), \quad a \leq t_0 < b_0, \quad t_0 \leq t \leq b_0, \end{aligned}$$

satisfy the following formula (analogous to the Lagrange one):

$$\begin{aligned}
& \int_{c_1}^{c_2} (d\mathbf{q}(t)f(t), z(t)) - \int_{c_1}^{c_2} (y(t), d\mathbf{q}(t)g(t)) \\
&= (iJy(c_2), z(c_2)) - (iJy(c_1), z(c_1)) \\
&+ \int_{c_1}^{c_2} (y(t), d\mathbf{p}_2(t)z(t)) - \int_{c_1}^{c_2} (d\mathbf{p}_1(t)y(t), z(t)) \\
&- \sum_{t \in \mathcal{S}_{\mathbf{p}_1} \cap \mathcal{S}_{\mathbf{p}_2} \cap [c_1, c_2)} (iJ\mathbf{p}_1(\{t\})y(t), \mathbf{p}_2(\{t\})z(t)) \\
&- \sum_{t \in \mathcal{S}_{\mathbf{q}} \cap \mathcal{S}_{\mathbf{p}_2} \cap [c_1, c_2)} (iJ\mathbf{q}(\{t\})f(t), \mathbf{p}_2(\{t\})z(t)) \\
&- \sum_{t \in \mathcal{S}_{\mathbf{p}_1} \cap \mathcal{S}_{\mathbf{q}} \cap [c_1, c_2)} (iJ\mathbf{p}_1(\{t\})y(t), \mathbf{q}(\{t\})g(t)) \\
&- \sum_{t \in \mathcal{S}_{\mathbf{q}} \cap [c_1, c_2)} (iJ\mathbf{q}(\{t\})f(t), \mathbf{q}(\{t\})g(t)), \quad t_0 \leq c_1 < c_2 \leq b_0. \quad (2.2)
\end{aligned}$$

Further we will assume that the measures  $\mathbf{p}$ ,  $\mathbf{m}$  have bounded variations,  $\mathbf{p}$  is self-adjoint and  $\mathbf{m}$  is non-negative. We consider the equation

$$y(t) = x_0 - iJ \int_a^t d\mathbf{p}(s)y(s) - iJ \int_a^t d\mathbf{m}(s)f(s), \quad (2.3)$$

where  $x_0 \in H$ ,  $f$  is integrable with respect to the measure  $\mathbf{m}$  on  $[a, b]$ ,  $a \leq t \leq b_0$ .

We construct a continuous measure  $\mathbf{p}_0$  from the measure  $\mathbf{p}$  in the following way. We set  $\mathbf{p}_0(\{t_k\}) = 0$  for  $t_k \in \mathcal{S}_{\mathbf{p}}$  and we set  $\mathbf{p}_0(\Delta) = \mathbf{p}(\Delta)$  for all Borel sets such that  $\Delta \cap \mathcal{S}_{\mathbf{p}} = \emptyset$ . Similarly, we construct a continuous measure  $\mathbf{m}_0$  from the measure  $\mathbf{m}$ . The measures  $\mathbf{p}_0$ ,  $\mathbf{m}_0$  are self-adjoint and the measure  $\mathbf{m}_0$  is non-negative. We replace  $\mathbf{p}$  by  $\mathbf{p}_0$  and  $\mathbf{m}$  by  $\mathbf{m}_0$  in (2.3). Then we obtain the equation

$$y(t) = x_0 - iJ \int_a^t d\mathbf{p}_0(s)y(s) - iJ \int_a^t d\mathbf{m}_0(s)f(s). \quad (2.4)$$

Equations (2.3) and (2.4) have unique solutions (see [8]).

By  $W$ , denote the operator solution of the equation

$$W(t)x_0 = x_0 - iJ \int_a^t d\mathbf{p}_0(s)W(s)x_0, \quad (2.5)$$

where  $x_0 \in H$ . Using Lemma 2.1, we get

$$W^*(t)JW(t) = J \quad (2.6)$$

by the standard method (see [11]). The functions  $t \rightarrow W(t)$  and  $t \rightarrow W^{-1}(t) = JW^*(t)J$  are continuous with respect to the uniform operator topology. Consequently, there exist constants  $\varepsilon_1 > 0$ ,  $\varepsilon_2 > 0$  such that the inequality

$$\varepsilon_1 \|x\|^2 \leq \|W(t)x\|^2 \leq \varepsilon_2 \|x\|^2 \quad (2.7)$$

holds for all  $x \in H, t \in [a, b_0]$ . The following Lemma 2.2 is established in [12] for the case of a continuous measure  $\mathbf{m}$ .

**Lemma 2.2.** *The function  $y$  is a solution of the equation*

$$y(t) = x_0 - iJ \int_a^t d\mathbf{p}_0(s)y(s)x - iJ \int_a^t d\mathbf{m}(s)f(s), \quad x_0 \in H, \quad a \leq t \leq b_0, \quad (2.8)$$

if and only if  $y$  has the form

$$y(t) = W(t)x_0 - W(t)iJ \int_a^t W^*(\xi)d\mathbf{m}(\xi)f(\xi). \quad (2.9)$$

*Proof.* Equation (2.8) has a unique solution (see [8]). It is enough to prove that if we substitute the function from the right-hand side of (2.9) instead of  $y$  in equation (2.8), then we get the identity. With this substitution, the right-hand side of (2.8) takes the form

$$\begin{aligned} x_0 - iJ \int_a^t d\mathbf{p}_0(s) \left( W(s)x_0 - W(s)iJ \int_a^s W^*(\xi)d\mathbf{m}(\xi)f(\xi) \right) - iJ \int_a^t d\mathbf{m}(s)f(s) \\ = x_0 - iJ \int_a^t d\mathbf{p}_0(s) W(s)x_0 \\ - J \int_a^t d\mathbf{p}_0(s) W(s)J \int_a^s W^*(\xi) d\mathbf{m}(\xi) f(\xi) - iJ \int_a^t d\mathbf{m}(s) f(s). \end{aligned} \quad (2.10)$$

We change the limits of integration in the third term of the right-hand side of (2.10). Then the third term takes the form

$$\begin{aligned} J \int_a^t d\mathbf{p}_0(s) W(s)J \int_a^s W^*(\xi)d\mathbf{m}(\xi) f(\xi) \\ = J \int_{[a,t)} \left( \int_{(\xi,t)} d\mathbf{p}_0(s) W(s) \right) JW^*(\xi) d\mathbf{m}(\xi) f(\xi) \\ = J \int_{[a,t)} \left( \int_{[\xi,t)} d\mathbf{p}_0(s) W(s) \right) JW^*(\xi) d\mathbf{m}(\xi) f(\xi) \\ - J \int_{[a,t)} \left( \int_{\{\xi\}} d\mathbf{p}_0(s) W(s) \right) JW^*(\xi) d\mathbf{m}(\xi) f(\xi). \end{aligned} \quad (2.11)$$

The last term in (2.11) is equal to zero since the measure  $\mathbf{p}_0$  is continuous. Using (2.5), we continue equality (2.10):

$$W(t)x_0 - \int_a^t J \left( \int_\xi^t d\mathbf{p}_0(s) W(s) \right) JW^*(\xi) d\mathbf{m}(\xi) f(\xi) - iJ \int_a^t d\mathbf{m}(s) f(s). \quad (2.12)$$

It follows from (2.5) that (2.12) is equal to

$$W(t)x_0 - \int_a^t i((W(t) - E) - (W(\xi) - E))JW^*(\xi)d\mathbf{m}(\xi) f(\xi) - iJ \int_a^t d\mathbf{m}(s) f(s)$$

$$\begin{aligned}
&= W(t)x_0 - i \int_a^t W(t)JW^*(\xi) d\mathbf{m}(\xi) f(\xi) \\
&\quad + i \int_a^t W(\xi)JW^*(\xi) d\mathbf{m}(\xi) f(\xi) - iJ \int_a^t d\mathbf{m}(s) f(s).
\end{aligned}$$

Taking into account (2.6), we continue the last equality

$$\begin{aligned}
W(t)x_0 - iW(t)J \int_a^t W^*(\xi) d\mathbf{m}(\xi) f(\xi) \\
+ iJ \int_a^t d\mathbf{m}(\xi) f(\xi) - iJ \int_a^t d\mathbf{m}(s) f(s) = y(t).
\end{aligned}$$

The lemma is proved.  $\square$

### 3. Linear relations generated by the integral equation

Let  $\mathbf{B}$  be a Hilbert space. A linear relation  $T$  is understood as a linear manifold  $T \subset \mathbf{B} \times \mathbf{B}$ . The terminology of the linear relations can be found, for example, in [1, 13]. In what follows we make use of the following notations:  $\{\cdot, \cdot\}$  is an ordered pair,  $\mathcal{D}(T)$  is the domain of  $T$ ,  $\mathcal{R}(T)$  is the range of  $T$ ,  $\ker T$  is the set of elements  $x \in \mathbf{B}$  such that  $\{x, 0\} \in T \subset \mathbf{B} \times \mathbf{B}$ . A relation  $T^*$  is called adjoint for  $T$  if  $T^*$  consists of all pairs  $\{y_1, y_2\}$  such that the equality  $(x_2, y_1) = (x_1, y_2)$  holds for all pairs  $\{x_1, x_2\} \in T$ . A linear relation  $T$  is called dissipative (accumulative, symmetric) if for any  $\{x, x'\} \in T$  we have  $\text{Im}(x', x) \geq 0$  (respectively,  $\text{Im}(x', x) \leq 0$ , or  $\text{Im}(x', x) = 0$ ). A dissipative (accumulative, symmetric) relation  $T$  is called maximal dissipative (accumulative, symmetric) if it has no dissipative (accumulative, symmetric) extensions  $T_1 \supset T$  such that  $T_1 \neq T$ . A symmetric relation is called self-adjoint if it is maximal dissipative and maximal accumulative at the same time. As it is known, a relation  $T$  is symmetric if and only if  $T \subset T^*$  and it is self-adjoint if and only if  $T = T^*$ . As linear operators are treated as linear relations, the notation  $\{x_1, x_2\} \in T$  is also used for the operator  $T$ . Since all considered relations are linear, we will often omit the word “linear”.

Let  $\mathbf{m}$  be a non-negative operator-valued measure defined on Borel sets  $\Delta \subset [a, b]$  that takes values in the set of linear bounded operators acting in the space  $H$ . The measure  $\mathbf{m}$  is assumed to have a bounded variation on  $[a, b]$ . We introduce the quasi-scalar product

$$(x, y)_{\mathbf{m}} = \int_a^{b_0} ((d\mathbf{m})x(t), y(t))$$

on a set of step-like functions with values in  $H$  defined on the segment  $[a, b_0]$ . Identifying with zero functions  $y$  obeying  $(y, y)_{\mathbf{m}} = 0$  and making the completion, we arrive at the Hilbert space denoted by  $L_2(H, d\mathbf{m}; a, b) = \mathfrak{H}$ . The elements of  $\mathfrak{H}$  are the classes of functions identified with respect to the norm  $\|y\|_{\mathbf{m}} = (y, y)_{\mathbf{m}}^{1/2}$ . In order not to complicate the terminology, the class of functions with

a representative  $y$  is indicated by the same symbol and we write  $y \in \mathfrak{H}$ . The equalities of the functions in  $\mathfrak{H}$  are understood as the equalities for associated equivalence classes.

Let us define a minimal relation  $L_0$  in the following way. The relation  $L_0$  consists of pairs  $\{\tilde{y}, \tilde{f}\} \in \mathfrak{H} \times \mathfrak{H}$  satisfying the condition: for each pair  $\{\tilde{y}, \tilde{f}\}$  there exists a pair  $\{y, f\}$  such that the pairs  $\{\tilde{y}, \tilde{f}\}$ ,  $\{y, f\}$  are identical in  $\mathfrak{H} \times \mathfrak{H}$  and  $\{y, f\}$  satisfies equation (2.3) and the equalities

$$y(a) = y(b_0) = y(\alpha) = 0, \quad \alpha \in \mathcal{S}_p; \quad \mathbf{m}(\{\beta\})f(\beta) = 0, \quad \beta \in \mathcal{S}_m. \quad (3.1)$$

In general, the relation  $L_0$  is not an operator since the function  $y$  may happen to be identified with zero in  $\mathfrak{H}$ , while  $f$  is non-zero. It follows from Lemma 2.1 that the relation  $L_0$  is symmetric. Further, without loss of generality it can be assumed that if a pair  $\{y, f\} \in L_0$ , then equalities (2.3) and (3.1) hold for this pair.

**Lemma 3.1.** *Equalities (2.3), (2.4), and (2.8) hold simultaneously for any pair  $\{y, f\} \in L_0$ .*

*Proof.* We denote  $\bar{\mathbf{p}} = \mathbf{p} - \mathbf{p}_0$ ,  $\bar{\mathbf{m}} = \mathbf{m} - \mathbf{m}_0$ . Then  $\bar{\mathbf{p}}(\{t_k\}) = \mathbf{p}(\{t_k\})$  for all  $t_k \in \mathcal{S}_p$  and  $\bar{\mathbf{p}}(\Delta) = 0$  for all Borel sets  $\Delta$  such that  $\Delta \cap \mathcal{S}_p = \emptyset$ . Similar equalities hold for the measure  $\bar{\mathbf{m}}$ . Using (2.3), we get

$$\begin{aligned} y(t) = x_0 - iJ \int_a^t d\mathbf{p}_0(s) y(s) - iJ \int_a^t d\bar{\mathbf{p}}(s) y(s) \\ - iJ \int_a^t d\bar{\mathbf{m}}(s) f(s) - iJ \int_a^t d\mathbf{m}_0(s) f(s). \end{aligned}$$

Now the desired statement follows from (3.1). The lemma is proved. □

**Corollary 3.2.** *If  $y \in \mathcal{D}(L_0)$ , then  $y$  is continuous and  $y(b) = 0$ .*

**Lemma 3.3.** *A pair  $\{\tilde{y}, \tilde{f}\} \in \mathfrak{H} \times \mathfrak{H}$  belongs to the relation  $L_0$  if and only if there exists a pair  $\{y, f\}$  such that the pairs  $\{\tilde{y}, \tilde{f}\}$ ,  $\{y, f\}$  are identical in  $\mathfrak{H} \times \mathfrak{H}$  and the equalities*

$$y(t) = -W(t) iJ \int_a^t W^*(s) d\mathbf{m}_0(s) f(s), \quad (3.2)$$

$$y(\alpha) = W(\alpha) iJ \int_a^\alpha W^*(s) d\mathbf{m}_0(s) f(s) = 0, \quad (3.3)$$

$$\mathbf{m}(\{\beta\})f(\beta) = 0 \quad (3.4)$$

hold, where  $\alpha \in \mathcal{S}_p \cup \{b_0\}$ ,  $\beta \in \mathcal{S}_m$ .

*Proof.* It follows from Lemmas 2.2 and 3.1 that equalities (3.2)–(3.4) hold together with equalities (2.3) and (3.1). By the definition of the relation  $L_0$ , a pair  $\{y, f\} \in L_0$  if and only if (2.3) and (3.1) hold. The lemma is proved. □

**Lemma 3.4.** *The relation  $L_0$  is closed.*

*Proof.* Suppose  $\{y_n, f_n\} \in L_0$ . Using (3.2)–(3.4), we obtain

$$y_n(t) = -W(t) iJ \int_a^t W^*(s) d\mathbf{m}_0(s) f_n(s), \tag{3.5}$$

$$y_n(\alpha) = W(\alpha) iJ \int_a^\alpha W^*(s) d\mathbf{m}_0(s) f_n(s) = 0, \quad \mathbf{m}(\{\beta\})f_n(\beta) = 0, \tag{3.6}$$

where  $\alpha \in \mathcal{S}_p \cup \{b_0\}$ ,  $\beta \in \mathcal{S}_m$ . Suppose that the sequences  $\{y_n\}$ ,  $\{f_n\}$  converge in  $\mathfrak{H}$  to  $y$ ,  $f$ , respectively. We note that if a sequence converges in  $\mathfrak{H} = L_2(H, d\mathbf{m}; a, b)$ , then this sequence converges in  $L_2(H, d\mathbf{m}_0; a, b)$ . Moreover,

$$\|f_n - f\|_{\mathfrak{H}}^2 \geq (\mathbf{m}(\{\beta\})(f_n(\beta) - f(\beta)), f_n(\beta) - f(\beta)) = (\mathbf{m}(\{\beta\})f(\beta), f(\beta)),$$

where  $\beta \in \mathcal{S}_m$ . Passing to the limit as  $n \rightarrow \infty$  in (3.5) and (3.6), we obtain equalities (3.2)–(3.4). It follows from Lemma 3.3 that the pair  $\{y, f\} \in L_0$ . The lemma is proved.  $\square$

**Corollary 3.5.** *The function  $f \in \mathfrak{H}$  belongs to the range  $\mathcal{R}(L_0)$  if and only if  $f$  satisfies the conditions*

$$\int_a^\alpha W^*(s) d\mathbf{m}_0(s) f(s) = 0, \quad \mathbf{m}(\{\beta\})f(\beta) = 0, \tag{3.7}$$

where  $\alpha \in \mathcal{S}_p \cup \{b_0\}$ ,  $\beta \in \mathcal{S}_m$ .

*Remark 3.6.* The first equality in (3.7) is equivalent to the following:

$$\int_{\alpha_1}^{\alpha_2} W^*(s) d\mathbf{m}_0(s) f(s) = 0, \quad \alpha_1, \alpha_2 \in \mathcal{S}_p \cup \{a\} \cup \{b_0\}. \tag{3.8}$$

*Remark 3.7.* It follows from Lemma 3.1, Corollary 3.2, and equality (3.4) that we can replace  $\mathbf{m}_0$  by  $\mathbf{m}$  and  $b_0$  by  $b$  in (3.2), (3.3), (3.7), and (3.8).

By  $\overline{\mathcal{S}_p}$ , denote the closure of the set  $\mathcal{S}_p$ .

**Lemma 3.8.** *Suppose  $\{y, f\} \in L_0$ . Then  $y(t) = 0$  for all  $t \in \overline{\mathcal{S}_p}$  and  $f(t) = 0$  for  $\mathbf{m}$ -almost all  $t \in \overline{\mathcal{S}_p} \cup \{a, b\}$ .*

*Proof.* It follows from Corollary 3.2 that the functions  $y \in \mathcal{D}(L_0)$  are continuous. Taking into account (3.1), we obtain  $y(t) = 0$  for  $t \in \overline{\mathcal{S}_p}$ . Using Corollary 3.5 and Remark 3.7, we get

$$\int_a^\alpha (d\mathbf{m}_0(s)f(s), W(s)x) = 0, \quad \mathbf{m}(\{\beta\})f(\beta) = 0$$

for all  $x \in H$  and for all  $\alpha \in \overline{\mathcal{S}_p} \cup \{b\}$ ,  $\beta \in \mathcal{S}_m$ . Hence equality (2.1) implies

$$\int_a^\alpha (\Psi_{\mathbf{m}_0}(s)f(s), W(s)x) d\rho_{\mathbf{m}_0}(s) = 0, \quad \mathbf{m}(\{\beta\})f(\beta) = 0. \tag{3.9}$$

We denote

$$\varphi_x(t) = (\Psi_{\mathbf{m}_0}(t)f(t), W(t)x), \quad \Phi_x(t) = \int_a^t \varphi_x(s) d\rho_{\mathbf{m}_0}(s).$$

The function  $\Phi_x$  is continuous. Hence, it follows from (3.9) that  $\Phi_x(t) = 0$  for all  $t \in \overline{\mathcal{S}_{\mathbf{p}}} \cup \{a, b\}$ . Therefore,  $\varphi_x(t) = 0$  for  $\rho_{\mathbf{m}_0}$ -almost all  $t \in \overline{\mathcal{S}_{\mathbf{p}}} \cup \{a, b\}$ .

Let  $\{x_n\}$  be a countable everywhere dense set in  $H$  and let  $\mathcal{X}_n$  be a set  $t \in \overline{\mathcal{S}_{\mathbf{p}}}$  such that  $\varphi_{x_n}(t) = 0$ . Then  $\varrho_{\mathbf{m}_0}(\mathcal{X}_n) = \varrho_{\mathbf{m}_0}(\overline{\mathcal{S}_{\mathbf{p}}})$ . We denote  $\mathcal{X} = \bigcap_n \mathcal{X}_n$ . Then  $\varrho_{\mathbf{m}_0}(\mathcal{X}) = \varrho_{\mathbf{m}_0}(\overline{\mathcal{S}_{\mathbf{p}}})$  and  $\varphi_{x_n}(t) = 0$  for all  $n$ . If a sequence  $\{z_n\}$  converges to  $z$  in  $H$ , then the sequence  $\{W(t)z_n\}$  converges to  $W(t)z$  for fixed  $t$ . Therefore,  $\varphi_x(t) = 0$  for all  $x \in H$  and for all  $t \in \mathcal{X}$ . The operator  $W(t)$  has a bounded inverse for all  $t$ . It follows that  $\Psi_{\mathbf{m}_0}(t)f(t) = 0$  for all  $t \in \mathcal{X}$ . Consequently,  $\Psi_{\mathbf{m}_0}(t)f(t) = 0$  for  $\rho_{\mathbf{m}_0}$ -almost all  $t \in \overline{\mathcal{S}_{\mathbf{p}}} \cup \{a, b\}$ . It follows from (2.1) that

$$\int_a^b (d\mathbf{m}_0(t)f(t), f(t)) = \int_a^b (\Psi_{\mathbf{m}_0}(t)f(t), f(t)) d\rho_{\mathbf{m}_0}(t) = 0.$$

Hence, using (3.4), we obtain  $f(t) = 0$  for  $\mathbf{m}$ -almost all  $t \in \overline{\mathcal{S}_{\mathbf{p}}} \cup \{a, b\}$ . The lemma is proved.  $\square$

By  $\mathfrak{H}_0$  (by  $\mathfrak{H}_1$ ), denote a subspace of functions that vanish on  $(a, b) \setminus \overline{\mathcal{S}_{\mathbf{p}}}$  (on  $\overline{\mathcal{S}_{\mathbf{p}}} \cup \{a, b\}$ , respectively) with respect to the norm in  $\mathfrak{H}$ . The subspaces  $\mathfrak{H}_0, \mathfrak{H}_1$  are orthogonal and  $\mathfrak{H} = \mathfrak{H}_0 \oplus \mathfrak{H}_1$ . We note that  $\mathfrak{H}_0 = \{0\}$  if and only if  $\mathbf{m}(\overline{\mathcal{S}_{\mathbf{p}}} \cup \{a, b\}) = 0$ .

We denote  $L_{10} = L_0 \cap (\mathfrak{H}_1 \times \mathfrak{H}_1)$ . Then  $\mathcal{D}(L_{10}) \subset \mathfrak{H}_1, \mathcal{R}(L_{10}) \subset \mathfrak{H}_1$ . It follows from Lemma 3.8 that

$$L_0^* = (\mathfrak{H}_0 \times \mathfrak{H}_0) \oplus L_{10}^*, \tag{3.10}$$

i.e., the relation  $L_0^*$  consists of all pairs  $\{y, f\} \in \mathfrak{H}$  of the form

$$\{y, f\} = \{u, v\} + \{z, g\} = \{u + z, v + g\},$$

where  $u, v \in \mathfrak{H}_0, \{z, g\} \in L_{10}^*$ .

The set  $\mathcal{T}_{\mathbf{p}} = (a, b) \setminus \overline{\mathcal{S}_{\mathbf{p}}}$  is open and it is the union of at most a countable number of disjoint open intervals, i.e.,  $\mathcal{T}_{\mathbf{p}} = \bigcup_{k=1}^{\mathbb{k}_1} \mathcal{J}_k, \mathcal{J}_k \cap \mathcal{J}_j = \emptyset$  for  $k \neq j$ , where  $\mathbb{k}_1$  is a natural number (equal to the number of intervals if this number is finite) or the symbol  $\infty$  (if the number of intervals is infinite). By  $\mathbb{J}$ , denote the set of these intervals  $\mathcal{J}_k$ . Note that the boundaries  $\alpha_k, \beta_k$  of any interval  $\mathcal{J}_k = (\alpha_k, \beta_k) \in \mathbb{J}$  belong to  $\overline{\mathcal{S}_{\mathbf{p}}} \cup \{a, b\}$ .

Further, let  $\chi_A$  denote the characteristic function of a set  $A$ . We denote

$$w_k(t) = \chi_{[\alpha_k, \beta_k)} W(t)W^{-1}(\alpha_k), \tag{3.11}$$

where  $(\alpha_k, \beta_k) = \mathcal{J}_k \in \mathbb{J}$ . Using (2.6), we get

$$w_k^*(t)Jw_k(t) = J, \quad \alpha_k \leq t < \beta_k. \tag{3.12}$$

**Lemma 3.9.** *Let  $g \in \mathfrak{H}$  and let the function  $G_k$  be given by the equality*

$$G_k(t) = -w_k(t) iJ \int_{\alpha_k}^t w_k^*(s) d\mathbf{m}(s) g(s),$$

where  $(\alpha_k, \beta_k) = \mathcal{J}_k \in \mathbb{J}$ . Then the pair  $\{G_k, g\} \in L_{10}^*$  if  $g$  vanishes outside  $[\alpha_k, \beta_k)$ .

*Proof.* Equalities (2.6) and (3.11) imply

$$G_k(t) = -\chi_{[\alpha_k, \beta_k)} W(t) iJ \int_{\alpha_k}^t W^*(s) d\mathbf{m}(s) g(s).$$

It follows from Lemma 2.2 that the function  $G_k$  is a solution of equation (2.8) on the segment  $[\alpha_k, \gamma]$ ,  $\gamma < \beta_k$  (for  $a = \alpha_k$ ,  $y = G_k$ ,  $f = g$ ,  $x_0 = 0$ ).

Suppose a pair  $\{y, f\} \in L_0$ . According to Lemma 3.1, the pair  $\{y, f\}$  satisfies equation (2.8) for  $x_0 = 0$ . Therefore we can apply formula (2.2) to the functions  $y, f, G_k, g$  for  $c_1 = \alpha_k$ ,  $c_2 = \gamma$ ,  $\mathbf{q} = \mathbf{m}$ ,  $\mathbf{p}_1 = \mathbf{p}_2 = \mathbf{p}_0$ . Since the measure  $\mathbf{p}_0$  is continuous, self-adjoint and (3.4) holds, we obtain

$$\int_{\alpha_k}^{\gamma} (g(s), d\mathbf{m}(s) y(s)) = \int_{\alpha_k}^{\gamma} (G_k(s), d\mathbf{m}(s) f(s)) + (iJG_k(\gamma), y(\gamma)). \tag{3.13}$$

The function  $y$  is continuous from the left and  $y(\beta_k) = 0$ . Hence, passing to the limit as  $\gamma \rightarrow \beta_k - 0$  in (3.13), we obtain

$$\int_{\alpha_k}^{\beta_k} (g(s), d\mathbf{m}(s) y(s)) = \int_{\alpha_k}^{\beta_k} (G_k(s), d\mathbf{m}(s) f(s)).$$

This implies the desired statement. The lemma is proved. □

Let  $\mathbb{M}$  be a set consisting of intervals  $\mathcal{J} \in \mathbb{J}$  and single-point sets  $\{\tau\}$ , where  $\tau \in \mathcal{S}_{\mathbf{m}} \setminus \overline{\mathcal{S}_{\mathbf{p}}}$ . The set  $\mathbb{M}$  is at most countable. We arrange the elements of  $\mathbb{M}$  in the form of a finite or infinite sequence and denote these elements by  $\mathcal{E}_k$ , where  $k$  is any natural number if the number of elements in  $\mathbb{M}$  is infinite, and  $1 \leq k \leq \mathbb{k}$  if the number of elements in  $\mathbb{M}$  is finite and equal to  $\mathbb{k}$ .

We will assign an operator function  $v_k$  to each element  $\mathcal{E}_k \in \mathbb{M}$  in the following way. If  $\mathcal{E}_k$  is the interval,  $\mathcal{E}_k = \mathcal{J}_k = (\alpha_k, \beta_k) \in \mathbb{J}$ , then

$$v_k(t) = \chi_{[\alpha_k, \beta_k) \setminus \mathcal{S}_{\mathbf{m}}} w_k(t). \tag{3.14}$$

If  $\mathcal{E}_k$  is a single-point set,  $\mathcal{E}_k = \{\tau_k\}$ ,  $\tau_k \in \mathcal{S}_{\mathbf{m}} \setminus (\overline{\mathcal{S}_{\mathbf{p}}} \cup \{a, b\})$ , and  $\tau_k \in \mathcal{J}_n = (\alpha_n, \beta_n)$ , then

$$v_k(t) = \chi_{\{\tau_k\}} w_n(\tau_k). \tag{3.15}$$

It follows from the definition of functions  $v_k$  that for each element  $x_1, x_2 \in H$  the functions  $v_k(\cdot)x_1, v_j(\cdot)x_2$  are orthogonal in  $\mathfrak{H}$  for  $k \neq j$ . Moreover,  $v_k(\cdot)x \in \mathfrak{H}_1$  for all  $x \in H$  and for all  $k$ .

**Lemma 3.10.** *The linear span of functions  $t \rightarrow v_k(t)\xi$ ,  $\xi \in H$ , is dense in  $\ker L_{10}^*$ . Here  $k \in \mathbb{N}$  if  $\mathbb{k} = \infty$ , and  $1 \leq k \leq \mathbb{k}$  if  $\mathbb{k}$  is finite.*

*Proof.* It follows from Corollary 3.5, Remark 3.7, and (3.10) that the range  $\mathcal{R}(L_{10})$  consists of all functions  $f \in \mathfrak{H}$  orthogonal to functions of the form  $v_k(\cdot)\xi$ , where  $\xi \in H$ . The equality  $\ker(L_{10}^*) \oplus \mathcal{R}(L_{10}) = \mathfrak{H}_1$  implies the desired assertion. The lemma is proved.  $\square$

Let  $Q_{k,0}$  be a set  $x \in H$  such that the functions  $t \rightarrow v_k(t)x$  are identical with zero in  $\mathfrak{H}$ . We put  $Q_k = H \ominus Q_{k,0}$ . On the linear space  $Q_k$ , we introduce a norm  $\|\cdot\|_-$  by the equality

$$\|\xi_k\|_- = \|v_k(\cdot)\xi_k\|_{\mathfrak{H}}, \quad \xi_k \in Q_k. \tag{3.16}$$

We note that if  $v_k$  has the form (3.14), then

$$\|\xi_k\|_- = \left( \int_{[\alpha_k, \beta_k] \setminus \mathcal{S}_m} (d\mathbf{m}(s) w_k(s)\xi_k, w_k(s)\xi_k) \right)^{1/2}, \quad \xi_k \in Q_k.$$

If  $v_k$  has the form (3.15), then

$$\|\xi_k\|_- = (\mathbf{m}(\{\tau_k\})w_n(\tau_k)\xi_k, w_n(\tau_k)\xi_k)^{1/2}, \quad \xi_k \in Q_k.$$

By  $Q_k^-$ , denote the completion of  $Q_k$  with respect to the norm (3.16). The norm (3.16) is generated by the scalar product  $(\xi_k, \eta_k)_- = (v_k(\cdot)\xi_k, v_k(\cdot)\eta_k)_{\mathfrak{H}}$ , where  $\xi_k, \eta_k \in Q_k$ . From the formula (2.1), in which the measure  $\mathbf{P}$  is replaced by  $\mathbf{m}$ , it follows that

$$\|\xi_k\|_- \leq \gamma \|\xi_k\|, \quad \xi_k \in Q_k, \tag{3.17}$$

where  $\gamma > 0$  is independent of  $\xi_k \in Q_k$ .

It follows from (3.17) that the space  $Q_k^-$  can be treated as a space with a negative norm with respect to  $Q_k$  [2, Chap. 1] and [13, Chap. 2]. By  $Q_k^+$ , we denote the associated space with a positive norm. The definition of spaces with positive and negative norms implies that  $Q_k^+ \subset Q_k$ . By  $(\cdot, \cdot)_+$  and  $\|\cdot\|_+$ , we denote the scalar product and the norm in  $Q_k^+$ , respectively.

Suppose that a sequence  $\{x_{kn}\}$ ,  $x_{kn} \in Q_k$ , converges in the space  $Q_k^-$  to  $x_0 \in Q_k^-$  as  $n \rightarrow \infty$ . Then a sequence  $\{v_k(\cdot)x_{kn}\}$  is fundamental in  $\mathfrak{H}$ . Therefore this sequence converges to some element  $x_0 \in \mathfrak{H}$ . We denote this element by  $v_k(\cdot)x_0$ .

Let  $\tilde{Q}_n^- = Q_1^- \times \dots \times Q_n^-$  ( $\tilde{Q}_n^+ = Q_1^+ \times \dots \times Q_n^+$ ) be the Cartesian product of the first  $n$  sets  $Q_k^-$  ( $Q_k^+$ , respectively) and let  $V_n(t) = (v_1(t), \dots, v_n(t))$  be the operator one-row matrix. It is convenient to treat elements from  $\tilde{Q}_n^-$  as one-column matrices, and to assume that  $V_n(t)\tilde{\xi}_n = \sum_{k=1}^n v_k(t)\xi_k$ , where we denote  $\tilde{\xi}_n = \text{col}(\xi_1, \dots, \xi_n) \in \tilde{Q}_n^-$ ,  $\xi_k \in Q_k^-$ .

Let  $\ker_k$  be a linear space of functions  $t \rightarrow v_k(t)\xi_k$ ,  $\xi_k \in Q_k^-$ . By (3.16), it follows that  $\ker_k$  is closed in  $\mathfrak{H}$ . The spaces  $\ker_k$  and  $\ker_j$  are orthogonal for  $k \neq j$ . We denote  $\mathcal{K}_n = \ker_1 \oplus \dots \oplus \ker_n$ . Obviously,  $\mathcal{K}_n \subset \mathcal{K}_m$  for  $n < m$ .

**Lemma 3.11.** *The set  $\bigcup_n \mathcal{K}_n$  is dense in  $\ker L_{10}^*$ .*

*Proof.* The required statement follows immediately from Lemma 3.10.  $\square$

By  $\mathcal{V}_n$ , denote the operator  $\tilde{\xi}_n \rightarrow V_n(\cdot)\tilde{\xi}_n$  ( $\tilde{\xi}_n \in \tilde{Q}_n^-$ ). The operator  $\mathcal{V}_n$  maps continuously and one-to-one  $\tilde{Q}_n^-$  onto  $\mathcal{K}_n \subset \mathfrak{H}_1 \subset \mathfrak{H}$ . Hence the adjoint operator  $\mathcal{V}_n^*$  maps  $\mathfrak{H}$  onto  $\tilde{Q}_n^+$  continuously. We find the form of the operator  $\mathcal{V}_n^*$ . For all  $\tilde{\xi}_n \in \tilde{Q}_n = Q_1 \times \dots \times Q_n$ ,  $f \in \mathfrak{H}$ , we have

$$(f, \mathcal{V}_n \tilde{\xi}_n)_{\mathfrak{H}} = \int_a^{b_0} (d\mathbf{m}(s) f(s), V_n(s)\tilde{\xi}_n) = \int_a^{b_0} (V_n^*(s) d\mathbf{m}(s) f(s), \tilde{\xi}_n) = (\mathcal{V}_n^* f, \tilde{\xi}_n).$$

Since  $\tilde{Q}_n$  is dense in  $\tilde{Q}_n^-$ , we obtain

$$\mathcal{V}_n^* f = \int_a^{b_0} V_n^*(s) d\mathbf{m}(s) f(s). \tag{3.18}$$

So we proved the following statement:

**Lemma 3.12.** *The operator  $\mathcal{V}_n$  maps continuously and one-to-one  $\tilde{Q}_n^-$  onto  $\mathcal{K}_n$ . The adjoint operator  $\mathcal{V}_n^*$  maps continuously  $\mathfrak{H}$  onto  $\tilde{Q}_n^+$  and acts by the formula (3.18). Moreover,  $\mathcal{V}_n^*$  maps one-to-one  $\mathcal{K}_n$  onto  $\tilde{Q}_n^+$ .*

Let  $\mathcal{Q}_-, \mathcal{Q}_+, \mathcal{Q}$  be linear spaces of sequences  $\tilde{\eta} = \{\eta_k\}$ ,  $\tilde{\varphi} = \{\varphi_k\}$ ,  $\tilde{\xi} = \{\xi_k\}$ , respectively, such that the series

$$\sum_{k=1}^{\mathbb{k}} \|\eta_k\|_-^2, \quad \sum_{k=1}^{\mathbb{k}} \|\varphi_k\|_+^2, \quad \sum_{k=1}^{\mathbb{k}} \|\xi_k\|^2$$

converge if  $\mathbb{k} = \infty$ , where  $\eta_k \in Q_k^-, \varphi_k \in Q_k^+, \xi_k \in Q_k$ . These spaces become Hilbert spaces if we introduce the scalar products by the formulas

$$\begin{aligned} (\tilde{\eta}, \tilde{\sigma})_- &= \sum_{k=1}^{\mathbb{k}} (\eta_k, \sigma_k)_-, & \tilde{\eta}, \tilde{\sigma} \in \mathcal{Q}_-, \\ (\tilde{\varphi}, \tilde{\psi})_+ &= \sum_{k=1}^{\mathbb{k}} (\varphi_k, \psi_k)_+, & \tilde{\varphi}, \tilde{\psi} \in \mathcal{Q}_+, \\ (\tilde{\xi}, \tilde{\zeta}) &= \sum_{k=1}^{\mathbb{k}} (\xi_k, \zeta_k), & \tilde{\xi}, \tilde{\zeta} \in \mathcal{Q}. \end{aligned}$$

In these spaces, the norms are defined by the equalities

$$\|\tilde{\eta}\|_-^2 = \sum_{k=1}^{\mathbb{k}} \|\eta_k\|_-^2, \quad \|\tilde{\varphi}\|_+^2 = \sum_{k=1}^{\mathbb{k}} \|\varphi_k\|_+^2, \quad \|\tilde{\xi}\|^2 = \sum_{k=1}^{\mathbb{k}} \|\xi_k\|^2.$$

The spaces  $\mathcal{Q}_+, \mathcal{Q}_-$  can be treated as spaces with positive and negative norms with respect to  $\mathcal{Q}$  (see [2, Chap. 1] and [13, Chap. 2]). So,  $\mathcal{Q}_+ \subset \mathcal{Q} \subset \mathcal{Q}_-$  and  $\varepsilon_1 \|\tilde{\varphi}\|_- \leq \|\tilde{\varphi}\| \leq \varepsilon_2 \|\tilde{\varphi}\|_+$ , where  $\tilde{\varphi} \in \mathcal{Q}_+, \varepsilon_1, \varepsilon_2 > 0$ . The ‘‘scalar product’’  $(\tilde{\eta}, \tilde{\varphi})$

is defined for all  $\tilde{\varphi} \in \mathcal{Q}_+$ ,  $\tilde{\eta} \in \mathcal{Q}_-$ . If  $\tilde{\eta} \in \mathcal{Q}$ , then  $(\tilde{\eta}, \tilde{\varphi})$  coincides with the scalar product in  $\mathcal{Q}$ .

Let  $\mathcal{M} \subset \mathcal{Q}_-$  be a set of sequences that vanish starting from a certain number (its own for each sequence). The set  $\mathcal{M}$  is dense in the space  $\mathcal{Q}_-$ . The operator  $\mathcal{V}_n$  is the restriction of  $\mathcal{V}_{n+1}$  to  $\tilde{\mathcal{Q}}_n^-$ . By  $\mathcal{V}'$ , denote an operator in  $\mathcal{M}$  such that  $\mathcal{V}'\tilde{\eta} = \mathcal{V}_n\tilde{\eta}_n$  for all  $n \in \mathbb{N}$ , where  $\tilde{\eta} = (\tilde{\eta}_n, 0, \dots)$ ,  $\tilde{\eta}_n \in \tilde{\mathcal{Q}}_n^-$ . It follows from (3.16) that  $\mathcal{V}'$  admits an extension by continuity to the space  $\mathcal{Q}_-$ . By  $\mathcal{V}$ , denote the extended operator. This operator maps continuously and one-to-one  $\mathcal{Q}_-$  onto  $\ker(L_{10}^*) \subset \mathfrak{H}_1 \subset \mathfrak{H}$ . Moreover, we denote  $\tilde{V}(t)\tilde{\eta} = (\mathcal{V}\tilde{\eta})(t)$ , where  $\tilde{\eta} = \{\eta_k\} \in \mathcal{Q}_-$ . Using (3.16), we get

$$(\mathcal{V}\tilde{\eta}, \mathcal{V}\tilde{\sigma})_{\mathfrak{H}} = (\tilde{\eta}, \tilde{\sigma})_-, \quad \tilde{\eta} = \{\eta_k\}, \quad \tilde{\sigma} = \{\sigma_k\}, \quad \tilde{\eta}, \tilde{\sigma} \in \mathcal{Q}_-. \tag{3.19}$$

The adjoint operator  $\mathcal{V}^*$  maps continuously  $\mathfrak{H}$  onto  $\mathcal{Q}_+$ . Let us find the form of  $\mathcal{V}^*$ . Suppose  $f \in \mathfrak{H}$ ,  $\tilde{\eta} \in \mathcal{M}$ ,  $\tilde{\eta} = \{\tilde{\eta}_n, 0, \dots\}$ . Then

$$(\tilde{\eta}, \mathcal{V}^*f) = (\mathcal{V}\tilde{\eta}, f)_{\mathfrak{H}} = \int_a^{b_0} (d\mathbf{m}(t) \tilde{V}(t)\tilde{\eta}, f(t)) = \int_a^{b_0} (\tilde{\eta}, \tilde{V}^*(t) d\mathbf{m}(t)f(t)).$$

Since  $\mathcal{V}^*f \in \mathcal{Q}_+$  and the set  $\mathcal{M}$  is dense in  $\mathcal{Q}_-$ , we get

$$\mathcal{V}^*f = \int_a^{b_0} \tilde{V}^*(t) d\mathbf{m}(t)f(t). \tag{3.20}$$

Taking into account Lemmas 3.11 and 3.12, we obtain the following statement.

**Lemma 3.13.** *The operator  $\mathcal{V}$  maps  $\mathcal{Q}_-$  onto  $\ker(L_{10}^*)$  continuously and one-to-one. A function  $z$  belongs to  $\ker(L_{10}^*)$  if and only if there exists an element  $\tilde{\eta} = \{\eta_k\} \in \mathcal{Q}_-$  such that  $z(t) = (\mathcal{V}\tilde{\eta})(t) = \tilde{V}(t)\tilde{\eta}$ . The operator  $\mathcal{V}^*$  maps  $\mathfrak{H}$  onto  $\mathcal{Q}_+$  continuously and acts by the formula (3.20), and  $\ker \mathcal{V}^* = \mathfrak{H}_0 \oplus \mathcal{R}(L_{10})$ . Moreover,  $\mathcal{V}^*$  maps  $\ker(L_{10}^*)$  onto  $\mathcal{Q}_+$  one-to-one.*

**Theorem 3.14.** *A pair  $\{\tilde{y}, \tilde{f}\} \in \mathfrak{H} \times \mathfrak{H}$  belongs to  $L_0^*$  if and only if there exists a pair  $\{y, f\}$ , the functions  $y_0, y'_0 \in \mathfrak{H}_0$ ,  $\hat{y}, \hat{f} \in \mathfrak{H}_1$  and an element  $\tilde{\eta} \in \mathcal{Q}_-$  such that the pairs  $\{\tilde{y}, \tilde{f}\}$ ,  $\{y, f\}$  are identical in  $\mathfrak{H} \times \mathfrak{H}$  and the equalities*

$$y = y_0 + \hat{y}, \quad f = y'_0 + \hat{f}, \quad \hat{y}(t) = \tilde{V}(t)\tilde{\eta} - \sum_{k=1}^{\mathbb{k}_1} w_k(t) iJ \int_a^t w_k^*(s) d\mathbf{m}(s) \hat{f}(s) \tag{3.21}$$

hold, where the series in (3.21) converges in  $\mathfrak{H}$ ,  $\mathbb{k}_1$  is the number of intervals  $\mathcal{J}_k \in \mathbb{J}$ .

*Proof.* The first two equalities in (3.21) follow from (3.10). Let us prove that the last equality in (3.21) holds. First we prove that if the functions  $\hat{y}, \hat{f}$  satisfy the third equality in (3.21), then the pair  $\{\hat{y}, \hat{f}\} \in L_{10}^*$ . If  $\mathbb{k}_1$  is finite, then this statement follows from Lemmas 3.9 and 3.13. We assume that  $\mathbb{k}_1 = \infty$ .

It follows from Lemma 3.13 that  $\mathcal{V}\tilde{\eta} \in \ker(L_{10}^*)$ . The function

$$\hat{y}_k(t) = -w_k(t) iJ \int_a^t w_k^*(s) d\mathbf{m}(s) \hat{f}(s)$$

$$= -w_k(t) iJ \int_{\alpha_k}^t w_k^*(s) \Psi_{\mathbf{m}}(s) \widehat{f}(s) d\rho_{\mathbf{m}}(s) \tag{3.22}$$

vanishes outside the interval  $[\alpha_k, \beta_k]$ . (Here  $\Psi_{\mathbf{m}}, \rho_{\mathbf{m}}$  are the functions from (2.1) in which the measure  $\mathbf{P}$  is replaced by  $\mathbf{m}$ .) We denote  $\widehat{f}_k(t) = \chi_{[\alpha_k, \beta_k]} \widehat{f}(t)$ . Using (2.1), (2.7), and (3.22), we get

$$\begin{aligned} \|\widehat{y}_k(t)\| &\leq \varepsilon_1 \|w_k(t)\| \int_{\alpha_k}^{\beta_k} \|w_k^*(s)\| \|\Psi_{\mathbf{m}}^{1/2}(s) \widehat{f}_k(s)\| d\rho_{\mathbf{m}}(s) \\ &\leq \varepsilon \left( \int_{\alpha_k}^{\beta_k} \|\Psi_{\mathbf{m}}^{1/2}(s) \widehat{f}_k(s)\|^2 d\rho_{\mathbf{m}}(s) \right)^{1/2} = \varepsilon \|\widehat{f}_k\|_{\mathfrak{H}}, \quad \varepsilon_1, \varepsilon > 0. \\ \|\widehat{y}_k\|_{\mathfrak{H}}^2 &= \int_{\alpha_k}^{\beta_k} (\Psi_{\mathbf{m}}(t) \widehat{y}_k(t), \widehat{y}_k(t)) d\rho_{\mathbf{m}}(t) \leq \varepsilon^2 \rho_{\mathbf{m}}([\alpha_k, \beta_k]) \|\widehat{f}_k\|_{\mathfrak{H}}^2. \end{aligned} \tag{3.23}$$

We denote

$$S_n(t) = \sum_{k=1}^n \widehat{y}_k(t)$$

and prove that the sequence  $\{S_n\}$  converges in  $\mathfrak{H}$ . From (3.23), we get

$$\|S_n\|_{\mathfrak{H}}^2 = \sum_{k=1}^n \|\widehat{y}_k\|_{\mathfrak{H}}^2 \leq \varepsilon^2 \sum_{k=1}^n \rho_{\mathbf{m}}([\alpha_k, \beta_k]) \|\widehat{f}_k\|_{\mathfrak{H}}^2 \leq \varepsilon^2 \rho_{\mathbf{m}}([a, b]) \|\widehat{f}\|_{\mathfrak{H}}^2.$$

Consequently, the sequence  $\{S_n\}$  converges to some function  $S \in \mathfrak{H}$  and

$$S(t) = - \sum_{k=1}^{\infty} w_k(t) iJ \int_a^t w_k^*(s) d\mathbf{m}(s) \widehat{f}(s), \quad \|S\|_{\mathfrak{H}} \leq \varepsilon_2 \|\widehat{f}\|_{\mathfrak{H}}, \quad \varepsilon_2 > 0. \tag{3.24}$$

It follows from Lemma 3.9 that

$$\left\{ S_n, \sum_{k=1}^n \widehat{f}_k \right\} \in L_{10}^*.$$

The relation  $L_{10}^*$  is closed. Therefore,  $\{S, \widehat{f}\} \in L_{10}^*$  and  $\{\widehat{y}, \widehat{f}\} \in L_{10}^*$ .

Now we assume that a pair  $\{\widehat{y}, \widehat{f}\} \in L_{10}^*$ . For the function  $\widehat{f}$ , we find a function  $S$  by the formula (3.24). Then  $\{S, \widehat{f}\} \in L_{10}^*$ . Hence  $\widehat{y} - S \in \ker L_{10}^*$ . By Lemma 3.13, it follows that there exists an element  $\widetilde{\eta} \in \mathcal{Q}_-$  such that  $\widehat{y} - S = \mathcal{V}\widetilde{\eta}$ . Therefore,  $\widehat{y}$  has the form (3.21). Now (3.10) implies the desired assertion. The theorem is proved.  $\square$

#### 4. The description of dissipative extensions of $L_0$

By  $\mathcal{L}_0$  (by  $\mathcal{L}_0^\perp$ ), denote the closure in  $\mathfrak{H}$  of the linear span of functions  $t \rightarrow v_k(t)\eta_k$ , where  $\eta_k \in \mathcal{Q}_k^-$  and  $v_k$  has the form (3.15) (form (3.14), respectively). The spaces  $\mathcal{L}_0$  and  $\mathcal{L}_0^\perp$  are orthogonal. Using Lemmas 3.10 and 3.13, we obtain  $\mathcal{L}_0 \oplus \mathcal{L}_0^\perp = \ker L_{10}^*$ . We put  $\mathcal{Q}_- = \mathcal{V}^{-1}\mathcal{L}_0$ ,  $\mathcal{Q}_-^\perp = \mathcal{V}^{-1}\mathcal{L}_0^\perp$ . By (3.19), it follows

that the spaces  $\Omega_-, \Omega_-^\perp$  are orthogonal in  $\mathcal{Q}_-$  and  $\mathcal{Q}_- = \Omega_- \oplus \Omega_-^\perp$ . We denote  $\mathcal{V}_0 = \mathcal{V}P, \mathcal{V}_0^\perp = \mathcal{V}(E - P)$ , where  $P$  is the orthogonal projection onto  $\Omega_-$  in  $\mathcal{Q}_-$ .

It follows from Lemma 3.13 that  $\mathcal{V}^*f$  ( $f \in \mathfrak{H}$ ) is an element of the space  $\mathcal{Q}_+ \subset \mathcal{Q}$ , i.e., a sequence with elements of the form

$$w_n^*(\tau_k)\mathbf{m}(\{\tau_k\})f(\tau_k), \int_a^{b_0} \chi_{[\alpha_k, \beta_k] \setminus \mathcal{S}_m} w_k^*(t) d\mathbf{m}(t) f(t) \tag{4.1}$$

(and possibly with zeros), where  $\tau_k \in (\mathcal{S}_m \setminus \overline{\mathcal{S}_p}) \cap \mathcal{J}_n; (\alpha_k, \beta_k) = \mathcal{J}_k; \mathcal{J}_n, \mathcal{J}_k \in \mathbb{J}$ . The element  $\mathcal{V}_0^*f$  is a sequence with elements of the first form in (4.1) (and possibly with zeros), and  $(\mathcal{V}_0^\perp)^*f$  is a sequence with elements of the second form in (4.1) (and possibly with zeros). Therefore,

$$(\mathcal{V}^*f, \mathcal{V}_0^*g) = (\mathcal{V}_0^*f, \mathcal{V}_0^*g), \quad f, g \in \mathfrak{H}. \tag{4.2}$$

Using (3.12), we obtain

$$(iJw_n^*(\tau_k)\mathbf{m}(\{\tau_k\})f(\tau_k), w_n^*(\tau_k)\mathbf{m}(\{\tau_k\})g(\tau_k)) = (iJ\mathbf{m}(\{\tau_k\})f(\tau_k), \mathbf{m}(\{\tau_k\})g(\tau_k)), \quad f, g \in \mathfrak{H}. \tag{4.3}$$

We denote  $\mathbf{H}_- = \mathfrak{H}_0 \times \mathcal{Q}_-, \mathbf{H}_+ = \mathfrak{H}_0 \times \mathcal{Q}_+$ . Suppose a pair  $\{\tilde{y}, \tilde{f}\} \in L_0^*$ . By Theorem 3.14, there exists a pair  $\{y, f\}$  such that the pairs  $\{\tilde{y}, \tilde{f}\}, \{y, f\}$  are identical in  $\mathfrak{H} \times \mathfrak{H}$  and the equalities

$$y = y_0 + \hat{y}, \quad f = y'_0 + \hat{f}, \quad \{\hat{y}, \hat{f}\} \in L_{10}^* \tag{4.4}$$

hold, where  $y_0, y'_0 \in \mathfrak{H}_0$  and  $\hat{y}$  has the form (3.21). With each pair  $\{y, f\}$  we associate a pair of boundary values  $\{Y, Y'\} \in \mathbf{H}_- \times \mathbf{H}_+$  by the formulas

$$Y = \{y_0, Y_{10}\} \in \mathbf{H}_- = \mathfrak{H}_0 \times \mathcal{Q}_-, \quad Y' = \{y'_0, Y'_{10}\} \in \mathbf{H}_+ = \mathfrak{H}_0 \times \mathcal{Q}_+, \tag{4.5}$$

where

$$Y_{10} = \tilde{\eta} - 2^{-1}i\tilde{J}\mathcal{V}^*\hat{f} + 2^{-1}i\tilde{J}\mathcal{V}_0^*\hat{f}, \quad Y'_{10} = \mathcal{V}^*\hat{f}, \tag{4.6}$$

$\tilde{J}$  is the operator in  $\mathcal{Q}$  acting as  $\tilde{J}\tilde{\xi} = \{J\xi_k\}, \tilde{\xi} = \{\xi_k\} \in \mathcal{Q}$ .

Let  $\Gamma$  denote the operator that takes each pair  $\{y, f\} \in L_0^*$  to the ordered pair  $\{Y, Y'\}$  of boundary values  $Y, Y'$ , i.e.,  $\Gamma\{y, f\} = \{Y, Y'\}$ . We put  $\Gamma_1\{y, f\} = Y, \Gamma_2\{y, f\} = Y'$ . It follows from Lemma 3.13 that if pairs  $\{\tilde{y}, \tilde{f}\}, \{y, f\}$  are identical in  $\mathfrak{H} \times \mathfrak{H}$ , then their boundary values coincide.

**Theorem 4.1.** *The range  $\mathcal{R}(\Gamma)$  of the operator  $\Gamma$  coincides with  $\mathbf{H}_- \times \mathbf{H}_+$  and “the Green formula”*

$$(f, z)_{\mathfrak{H}} - (y, g)_{\mathfrak{H}} = (Y', Z) - (Y, Z') \tag{4.7}$$

holds, where  $\{y, f\}, \{z, g\} \in L_0^*, \Gamma\{y, f\} = \{Y, Y'\}, \Gamma\{z, g\} = \{Z, Z'\}$ .

*Proof.* The equality  $\mathcal{R}(\Gamma) = \mathbf{H}_- \times \mathbf{H}_+$  follows from Lemma 3.13 and the formulas (3.10), (4.5), (4.6). Let us prove (4.7). Suppose that a pair  $\{y, f\}$  has the form (3.21) and a pair  $\{z, g\}$  has the form

$$z = z_0 + \widehat{z}, \quad g = z'_0 + \widehat{g}, \quad \{\widehat{z}, \widehat{g}\} \in L_{10}^*$$

where  $z_0, z'_0 \in \mathfrak{H}_0$ ,

$$\widehat{z}(t) = \widetilde{V}(t)\widetilde{\zeta} - \sum_{k=1}^{k_1} w_k(t) iJ \int_a^t w_k^*(s) d\mathbf{m}(s) \widehat{g}(s), \quad \widetilde{\zeta} \in \mathcal{Q}_-, \quad \widehat{g} \in \mathfrak{H}_1. \quad (4.8)$$

Then

$$(f, z)_{\mathfrak{H}} - (y, g)_{\mathfrak{H}} = (y'_0, z_0)_{\mathfrak{H}} - (y_0, z'_0)_{\mathfrak{H}} + (\widehat{f}, \widehat{z})_{\mathfrak{H}} - (\widehat{y}, \widehat{g})_{\mathfrak{H}}.$$

Thus, it is enough to prove the equality

$$(\widehat{f}, \widehat{z})_{\mathfrak{H}} - (\widehat{y}, \widehat{g})_{\mathfrak{H}} = (Y'_{10}, Z_{10}) - (Y_{10}, Z'_{10}). \quad (4.9)$$

Using (4.6), we get

$$(\widehat{f}, \mathcal{V}\widetilde{\zeta})_{\mathfrak{H}} = (\mathcal{V}^*\widehat{f}, \widetilde{\zeta}) = (\mathcal{V}^*\widehat{f}, Z_{10} + 2^{-1}i\widetilde{J}\mathcal{V}^*\widehat{g} - 2^{-1}i\widetilde{J}\mathcal{V}_0^*\widehat{g}), \quad (4.10)$$

$$(\mathcal{V}\widetilde{\eta}, \widehat{g})_{\mathfrak{H}} = (\widetilde{\eta}, \mathcal{V}^*g) = (Y_{10} + 2^{-1}i\widetilde{J}\mathcal{V}^*\widehat{f} - 2^{-1}i\widetilde{J}\mathcal{V}_0^*\widehat{f}, \mathcal{V}^*\widehat{g}). \quad (4.11)$$

In (3.21) and (4.8), we denote

$$\begin{aligned} \widetilde{F}(t) &= - \sum_{k=1}^{k_1} w_k(t) iJ \int_a^t w_k^*(s) d\mathbf{m}(s) \widehat{f}(s), \\ \widetilde{G}(t) &= - \sum_{k=1}^{k_1} w_k(t) iJ \int_a^t w_k^*(s) d\mathbf{m}(s) \widehat{g}(s). \end{aligned}$$

We define the functions  $F_k, G_k$  by the equalities

$$F_k(t) = -w_k(t) iJ \int_{\alpha_k}^t w_k^*(s) d\mathbf{m}(s) \widehat{f}(s), \quad G_k(t) = -w_k(t) iJ \int_{\alpha_k}^t w_k^*(s) d\mathbf{m}(s) \widehat{g}(s).$$

It follows from Lemma 2.2 that the functions  $F_k, G_k$  are the solutions of equation (2.8) on  $[\alpha_k, \beta_k]$  for  $x_0 = 0$  ( $G_k$  is the solution if  $f$  is replaced by  $g$  in (2.8)). Using (3.12) and Lemma 2.1, for  $\mathbf{p}_1 = \mathbf{p}_2 = \mathbf{p}_0, \mathbf{q} = \mathbf{m}, c_1 = \alpha_k, c_2 = \beta < \beta_k$ , we obtain

$$\begin{aligned} &\int_{\alpha_k}^{\beta} (\widehat{f}(s), d\mathbf{m}(s) G_k(s)) - \int_{\alpha_k}^{\beta} (F_k(s), d\mathbf{m}(s) \widehat{g}(s)) \\ &= \left( iJ w_k(\beta) iJ \int_{\alpha_k}^{\beta} w_k^*(s) d\mathbf{m}(s) \widehat{f}(s), w_k(\beta) iJ \int_{\alpha_k}^{\beta} w_k^*(s) d\mathbf{m}(s) \widehat{g}(s) \right) \\ &\quad - \sum_{\tau \in \mathcal{S}_{\mathbf{m}} \cap [\alpha_k, \beta]} (iJ \mathbf{m}(\{\tau\}) \widehat{f}(\tau), \mathbf{m}(\{\tau\}) \widehat{g}(\tau)) \end{aligned}$$

$$\begin{aligned}
 &= \left( iJ \int_{\alpha_k}^{\beta} w_k^*(s) d\mathbf{m}(s) \widehat{f}(s), \int_{\alpha_k}^{\beta} w_k^*(s) d\mathbf{m}(s) \widehat{g}(s) \right) \\
 &\quad - \sum_{\tau \in \mathcal{S}_{\mathbf{m}} \cap [\alpha_k, \beta)} (iJ\mathbf{m}(\{\tau\})\widehat{f}(\tau), \mathbf{m}(\{\tau\})\widehat{g}(\tau)). \tag{4.12}
 \end{aligned}$$

Passing to the limit as  $\beta \rightarrow \beta_k - 0$  in (4.12), we obtain that (4.12) will remain true if  $\beta$  is replaced by  $\beta_k$ . Therefore,

$$\begin{aligned}
 &\int_{\alpha_k}^{\beta_k} (\widehat{f}(s), d\mathbf{m}(s) G_k(s)) - \int_{\alpha_k}^{\beta_k} (F_k(s), d\mathbf{m}(s) \widehat{g}(s)) \\
 &= \left( iJ \int_{\alpha_k}^{\beta_k} w_k^*(s) d\mathbf{m}(s) \widehat{f}(s), \int_{\alpha_k}^{\beta_k} w_k^*(s) d\mathbf{m}(s) \widehat{g}(s) \right) \\
 &\quad - \sum_{\tau \in \mathcal{S}_{\mathbf{m}} \cap [\alpha_k, \beta_k)} (iJ\mathbf{m}(\{\tau\})\widehat{f}(\tau), \mathbf{m}(\{\tau\})\widehat{g}(\tau)).
 \end{aligned}$$

Taking into account (3.20), (4.1), and (4.3), we get

$$(\widehat{f}, \widetilde{G})_{\mathfrak{H}} - (\widetilde{F}, \widehat{g})_{\mathfrak{H}} = (i\widetilde{J}\mathcal{V}^*\widehat{f}, \mathcal{V}^*\widehat{g}) - (i\widetilde{J}\mathcal{V}_0^*\widehat{f}, \mathcal{V}_0^*\widehat{g}).$$

Then equalities (4.10) and (4.11) imply

$$\begin{aligned}
 (\widehat{f}, \widehat{z})_{\mathfrak{H}} - (\widehat{y}, \widehat{g})_{\mathfrak{H}} &= (\mathcal{V}^*\widehat{f}, Z_{10}) - 2^{-1}(i\widetilde{J}\mathcal{V}^*\widehat{f}, \mathcal{V}^*\widehat{g}) + 2^{-1}(i\widetilde{J}\mathcal{V}^*\widehat{f}, \mathcal{V}_0^*\widehat{g}) \\
 &\quad - (Y_{10}, \mathcal{V}^*\widehat{g}) - 2^{-1}(i\widetilde{J}\mathcal{V}^*\widehat{f}, \mathcal{V}^*\widehat{g}) + 2^{-1}(i\widetilde{J}\mathcal{V}_0^*\widehat{f}, \mathcal{V}^*\widehat{g}) \\
 &\quad + (i\widetilde{J}\mathcal{V}^*\widehat{f}, \mathcal{V}^*\widehat{g}) - (i\widetilde{J}\mathcal{V}_0^*\widehat{f}, \mathcal{V}_0^*\widehat{g}).
 \end{aligned}$$

Now, using (4.2) and (4.6), we obtain (4.9). The theorem is proved. □

From the theory of spaces with positive and negative norms (see [2, Chap. 1] and [13, Chap. 2]), it follows that there exist isometric operators  $\delta_- : \mathcal{Q}_- \rightarrow \mathcal{Q}$ ,  $\delta_+ : \mathcal{Q}_+ \rightarrow \mathcal{Q}$  such that the equality  $(\widetilde{\eta}, \widetilde{\varphi}) = (\delta_- \widetilde{\eta}, \delta_+ \widetilde{\varphi})$  holds for all  $\widetilde{\eta} \in \mathcal{Q}_-$ ,  $\widetilde{\varphi} \in \mathcal{Q}_+$ . We denote  $\mathcal{H} = \mathfrak{H}_0 \times \mathcal{Q}$ . Suppose  $\{\widetilde{y}, \widehat{f}\} \in L_0^*$ . According to Theorem 3.14, there exists a pair  $\{y, f\}$  such that the pairs  $\{\widetilde{y}, \widehat{f}\}$ ,  $\{y, f\}$  are identical in  $\mathfrak{H} \times \mathfrak{H}$  and equalities (4.4) hold. To each pair  $\{y, f\}$  assign a pair of boundary values  $\gamma\{y, f\} = \{\mathcal{Y}, \mathcal{Y}'\} \in \mathcal{H} \times \mathcal{H}$  by the formulas

$$\mathcal{Y} = \gamma_1\{y, f\} = \{y_0, \delta_- Y_{10}\}, \quad \mathcal{Y}' = \gamma_2\{y, f\} = \{y'_0, \delta_+ Y'_{10}\}.$$

By Theorem 4.1, it follows that the operator  $\gamma$  maps  $L_0^*$  onto  $\mathcal{H} \times \mathcal{H}$  and the equality

$$(f, z)_{\mathfrak{H}} - (y, g)_{\mathfrak{H}} = (\mathcal{Y}', \mathcal{Z}) - (\mathcal{Y}, \mathcal{Z}') \tag{4.13}$$

holds, where  $\{y, f\}, \{z, g\} \in L_0^*$ ,  $\gamma\{y, f\} = \{\mathcal{Y}, \mathcal{Y}'\}$ ,  $\gamma\{z, g\} = \{\mathcal{Z}, \mathcal{Z}'\}$ . This implies that the ordered triple  $(\mathcal{H}, \gamma_1, \gamma_2)$  is a space of boundary values (a boundary triplet in another terminology) for  $L_0$  in the sense of papers [4, 5, 16] (see also [13, Chap. 3]).

Let  $\theta$  be a linear relation,  $\theta \subset \mathcal{H} \times \mathcal{H}$ . By  $L_\theta$ , denote a linear relation such that  $L_0 \subset L_\theta \subset L_0^*$  and  $\gamma L_\theta = \theta$ . It follows from (4.13) that both relations  $L_\theta$

and  $\theta$  are maximal dissipative (or maximal accumulative, or maximal symmetric, or self-adjoint). From here, taking into account the description of self-adjoint relations (see [20]), of dissipative relations (see [14]), we obtain the following assertion.

**Theorem 4.2.** *If  $U$  is a contraction on  $\mathfrak{H}$ , then the restriction of the relation  $L_0^*$  to the set of pairs  $\{y, f\} \in L_0^*$  satisfying the condition*

$$(U - E)\Gamma_2 f + (U + E)\Gamma_1 f = 0 \quad (4.14)$$

or

$$(U - E)\Gamma_2 f - (U + E)\Gamma_1 f = 0 \quad (4.15)$$

is a maximal dissipative, respectively, maximal accumulative extension of  $L_0$ . Conversely, any maximal dissipative (maximal accumulative) extension of  $L_0$  is the restriction of  $L_0^*$  to the set of pairs  $\{y, f\} \in L_0^*$  satisfying (4.14) (or (4.15)), where a contraction  $U$  is uniquely determined by an extension. The maximal symmetric extensions of the relation  $L_0$  on  $\mathfrak{H}$  are described by the conditions (4.14), (or (4.15)), where  $U$  is an isometric operator. These conditions define a self-adjoint extension if  $U$  is unitary.

Let us consider some examples.

*Example 4.3.* Suppose  $\mathbf{p} = \mathbf{p}_0$  is a continuous measure,  $\mathbf{m} = \mu$  is the usual Lebesgue measure on  $[a, b]$  (i.e.,  $\mu([\alpha, \beta]) = \beta - \alpha$ , where  $a \leq \alpha < \beta \leq b$  (we write  $ds$  instead of  $d\mu(s)$ )). In this case,  $L_0, L_0^*$  are operators,  $\mathbb{k}_1 = \mathbb{k} = 1$ ,  $\mathfrak{H}_0 = \{0\}$ ,  $Q_{1,0} = \{0\}$ ,  $Q_1 = H = \mathcal{Q}_- = \mathcal{Q}_+$ . Equality (3.21) has the form

$$y(t) = W(t)\eta - W(t) iJ \int_a^t W^*(s)f(s) ds, \quad f = L_0^* y, \quad \eta \in H.$$

By direct calculations, we obtain

$$Y = 2^{-1}(y(a) + W^{-1}(b)y(b)); \quad Y' = iJ(W^{-1}(b)y(b) - y(a)). \quad (4.16)$$

Now we assume that the measures  $\mathbf{p}, \mathbf{m}$  are continuous. Generally, then  $L_0, L_0^*$  are not operators. In this case,  $\mathbb{k}_1 = \mathbb{k} = 1$ ,  $\mathfrak{H}_0 = \{0\}$ . In general,  $Q_1 \neq H$ ,  $Q_1 \neq Q_1^-$ . If a pair  $\{y, f\} \in L_0^*$  is such that  $y(a) \in Q_1$ , then equalities (4.16) hold.

Suppose that  $\mathbf{m} = \mu$  and the set  $\mathcal{S}_{\mathbf{p}}$  of single-point atoms of the measure  $\mathbf{p}$  can be arranged as an increasing sequence converging to  $b$ . For this case the space of boundary values was constructed in [11].

*Example 4.4.* Suppose that  $\mathcal{S}_{\mathbf{m}} \neq \emptyset$  and  $\mathbf{m} = \mu + \overline{\mathbf{m}}$ , where  $\mu = \mathbf{m}_0$  is the usual Lebesgue measure on  $[a, b]$  and  $\mu(\Delta) = \mathbf{m}(\Delta)$  for all Borel sets such that  $\Delta \cap \mathcal{S}_{\mathbf{m}} = \emptyset$ . So,  $\mathcal{S}_{\mathbf{m}} = \mathcal{S}_{\overline{\mathbf{m}}}$  and  $\mathbf{m}(\{\beta\}) = \overline{\mathbf{m}}(\{\beta\})$  for all  $\beta \in \mathcal{S}_{\mathbf{m}}$ . We denote  $\widehat{Q}_{k,0} = \ker \mathbf{m}(\{\tau_k\})$ ,  $\widehat{Q}_k = H \ominus \widehat{Q}_{k,0}$ , where  $\tau_k \in \mathcal{S}_{\mathbf{m}}$ . Let  $\mathbf{m}_k$  be the restriction

of the operator  $\mathbf{m}(\{\tau_k\})$  to  $\widehat{Q}_k$ . The operator  $\mathbf{m}_k$  is self-adjoint and  $\mathcal{R}(\mathbf{m}_k) \subset \widehat{Q}_k$ . By  $\widehat{Q}_k^-$ , denote the completion of  $\widehat{Q}_k$  with respect to the norm  $\|\xi\|_- = (\mathbf{m}_k \xi, \xi)^{1/2}$ , where  $\xi \in \widehat{Q}_k$ . Let  $\widehat{Q}_-$  be the linear space of sequences  $\tilde{\eta} = \{\eta_k\}$  such that the series  $\sum_{k=1}^{\infty} \|\eta_k\|_-^2$  converges if  $\mathbb{k}_2 = \infty$ , where  $\mathbb{k}_2$  is the number of elements in  $\mathcal{S}_m$ . Then  $\mathfrak{H} = L_2(H; a, b) \oplus \widehat{Q}_-$ .

Suppose  $\mathbf{p} = 0$  and  $a, b \notin \mathcal{S}_m$ . (The case of an arbitrary continuous measure  $\mathbf{p}$  can be considered in a similar way.) If  $\mathbf{p} = 0$ , then  $\mathfrak{H}_0 = \{0\}$ ,  $W(t) = E$ , and  $\mathcal{Q}_- = H \oplus \widehat{Q}_-$ . It follows from Lemma 3.3 and (3.1) that a pair  $\{y, f\} \in L_0$  if and only if

$$y(t) = -iJ \int_a^t f(s) ds, \quad y(b) = 0, \quad \mathbf{m}(\beta)f(\beta) = 0, \quad \beta \in \mathcal{S}_m.$$

Using Theorem 3.14, we obtain that a pair  $\{y, f\} \in L_0^*$  if and only if

$$y(t) = \eta_0 + \sum_{\tau_k \leq t} \chi_{\{\tau_k\}} \eta_k - iJ \int_a^t d\mathbf{m}(s) f(s), \tag{4.17}$$

where  $\eta_0 \in H$ ,  $\tau_k \in \mathcal{S}_m$ ,  $\eta_k \in \widehat{Q}_k^-$ , and the sequence  $\tilde{\eta} = \{\eta_0, \eta_k\}$  belongs to  $\mathcal{Q}_-$  (here  $k \in \mathbb{N}$  if  $\mathbb{k}_2 = \infty$ , and  $1 \leq k \leq \mathbb{k}_2$  if  $\mathbb{k}_2$  is finite).

It follows from (4.5), (4.6), and (4.1) that the boundary values  $Y, Y'$  are the sequence of the form

$$Y = \left\{ \eta_0 - 2^{-1}iJ \int_a^b f(s) ds, \eta_k \right\},$$

$$Y' = \left\{ \int_a^b f(s) ds, \mathbf{m}(\{\tau_k\})f(\tau_k) \right\}, \quad k = 1, 2, \dots$$

Suppose that the set  $\mathcal{S}_m$  of single-point atoms  $\tau_k$  of measure  $\mathbf{m}$  can be arranged as an increasing sequence;  $\tau_1 < \tau_2 < \dots$ . In this case, we find  $\eta_0, \eta_k$ . Using (4.17), we get

$$y(t) = \eta_0 + \sum_{\tau_k \leq t} \chi_{\{\tau_k\}} \eta_k - iJ \int_a^t f(s) ds - iJ \sum_{\tau_k < t} \mathbf{m}(\{\tau_k\})f(\tau_k). \tag{4.18}$$

From (4.18), by direct calculations we obtain

$$\eta_0 = y(a),$$

$$\eta_1 = y(\tau_1) - y(a) + iJ \int_a^{\tau_1} f(s) ds,$$

$$\eta_k = y(\tau_k) - y(\tau_{k-1}) + iJ \int_{\tau_{k-1}}^{\tau_k} f(s) ds + iJ \mathbf{m}(\{\tau_{k-1}\})f(\tau_{k-1}).$$

Thus, the boundary values  $Y, Y'$  are expressed through the values of the functions  $y, f$  and the integrals of  $f$ .

**Acknowledgment.** The author thanks the reviewer for useful comments.

## References

- [1] A.G. Baskakov, *Analysis of linear differential equations by methods of the spectral theory of difference operators and linear relations*, Uspekhi Mat. Nauk **68** (2013), No. 1, 77–128; Engl. transl.: Russian Math. Surveys **68** (2013), No. 1, 69–116.
- [2] Yu.M. Berezanski, *Expansions in Eigenfunctions of Selfadjoint Operators*, Naukova Dumka, Kiev, 1965; Engl. transl.: Amer. Math. Soc., Providence, RI, 1968.
- [3] J. Behrndt, S. Hassi, H. Snoo, and R. Wietsma, *Square-integrable solutions and Weil functions for singular canonical systems*, Math. Nachr. **284** (2011), No. 11–12, 1334–1384.
- [4] V.M. Bruk, *On a class of boundary value problems with spectral parameter in the boundary condition*, Mat. Sbornik **100** (1976), No. 2, 210–216; Engl. transl.: Math. USSR-Sbornik **29** (1976), No. 2, 186–192.
- [5] V.M. Bruk, *Extensions of symmetric relations*, Mat. Zametki **22** (1977), No. 6, 825–834; Engl. transl.: Mathematical Notes **22** (1977), No. 6, 953–958.
- [6] V.M. Bruk, *On a number of linearly independent square-integrable solutions of systems of differential equations*, Functional Analysis **5** (1975), Uljanovsk, 25–33.
- [7] V.M. Bruk, *Linear relations in a space of vector functions*, Mat. Zametki **24** (1978), No. 4, 499–511; Engl. transl.: Mathematical Notes, **24** (1978), No. 4, 767–773.
- [8] V.M. Bruk, *Boundary value problems for integral equations with operator measures*, Probl. Anal. Issues Anal. **6(24)** (2017), No. 1, 19–40.
- [9] V.M. Bruk, *On self-adjoint extensions of operators generated by integral equations*, Taurida Journal of Computer Science Theory and Mathematics (2017), No. 1(34), 17–31.
- [10] V.M. Bruk, *On the characteristic operator of an integral equation with a Nevanlinna measure in the infinite-dimensional case*, Zh. Mat. Fiz. Anal. Geom. **10** (2014), No. 2, 163–188.
- [11] V.M. Bruk, *Generalized Resolvents of operators generated by integral equations*, Probl. Anal. Issues Anal, **7(25)** (2018), No. 2, 20–38.
- [12] V.M. Bruk, *On self-adjoint extensions of relations generated by integral equations*, Taurida Journal of Computer Science Theory and Mathematics (2019), No. 1(42), 43–61.
- [13] V.I. Gorbachuk and M.L. Gorbachuk, *Boundary Value Problems for Differential-Operator Equations*, Naukova Dumka, Kiev, 1984; Engl. transl.: Kluwer Acad. Publ., Dordrecht-Boston-London, 1991.
- [14] M.L. Gorbachuk, A.N. Kochubei, and M.A. Rybak, *Dissipative extensions of differential operators in a space of vector-functions*, Dokl. AN USSR **205** (1971), No. 5, 1029–1032; Engl. transl.: Soviet Math. Dokl. **13** (1972), No. 1, 1063–1067.
- [15] V. Khrabustovskyi, *Analogs of generalized resolvents for relations generated by a pair of differential operator expressions one of which depends on spectral parameter in nonlinear manner*, Zh. Mat. Fiz. Anal. Geom. **9** (2013), No. 4, 496–535.
- [16] A.N. Kochubei, *Extensions of symmetric operators and symmetric binary relations*, Mat. Zametki **17** (1975), No. 1, 41–48; Engl. transl.: Mathematical Notes, **17** (1975), No. 1, 25–28.

- [17] M.G. Krein, *The theory of self-adjoint extensions of semi-bounded Hermitian transformations and its applications*, Mat. Sbornik **20** (1947), No. 3, 431–495; **21** (1947), No. 3, 365–404.
- [18] M.A. Naimark, *Linear Differential Operators*, Nauka, Moscow, 1969; Engl. transl.: George G. Harrar and Company, LTD, London-Toronto-Wellington-Sidney, 1969.
- [19] B.C. Orcutt, *Canonical Differential Equations*, Ph. D. Thesis, University of Virginia, 1969.
- [20] F.S. Rofe-Beketov, *Self-adjoint extensions of differential operators in a space of vector functions*, Dokl. AN USSR **184** (1969), No. 5, 1034–1037; Engl. transl.: Soviet Math. Dokl. **10** (1969), No. 1, 188–192.
- [21] F.S. Rofe-Beketov and A.M. Kholkin, *Spectral Analysis of Differential Operators*, World Scientific Monograph Series in Mathematics, **7**, Singapore, 2005.

Received October 26, 2019, revised December 10, 2019.

Vladislav M. Bruk,

*Saratov State Technical University, 77 Politekhnicheskaya str., Saratov 410054, Russia,*

E-mail: [vladislavbruk@mail.ru](mailto:vladislavbruk@mail.ru)

**Дисипативні розширення лінійних відношень,  
породжених інтегральними рівняннями з  
операторними мірами**

Vladislav M. Bruk

У статті визначено мінімальне відношення  $L_0$ , яке породжене інтегральним рівнянням з операторними мірами, і надано опис спряженого відношення  $L_0^*$ . Для цього мінімального відношення побудовано простір граничних значень (гранична трійка), що задовольняє абстрактну “формулу Гріна”, і одержано опис максимального дисипативного (аккумулятивного) відношення, а також самоспряжених розширень мінімального відношення.

*Ключові слова:* гільбертів простір, лінійне відношення, інтегральне рівняння, дисипативне розширення, самоспряжене розширення, граничне значення, операторна міра